

RANKIN TRIPLE PRODUCTS AND QUANTUM CHAOS

THOMAS C. WATSON

A DISSERTATION
PRESENTED TO THE FACULTY
OF PRINCETON UNIVERSITY
IN CANDIDACY FOR THE DEGREE
OF DOCTOR OF PHILOSOPHY

RECOMMENDED FOR ACCEPTANCE
BY THE DEPARTMENT OF
MATHEMATICS

JUNE 2002

REVISED 10/22/2008

© Copyright by Thomas C. Watson, 2002.

All Rights Reserved

Abstract

In this paper we demonstrate the chaotic nature of certain special arithmetic quantum dynamical systems, using machinery from analytic number theory.

Consider the quantized geodesic flow on a finite-volume hyperbolic surface $\Gamma \backslash \mathfrak{H}$, with $\Gamma \subset \mathrm{SL}_2 \mathbb{R}$ consisting of the norm-1 units of an Eichler order in an indefinite quaternion algebra B over $\mathbb{Q}$. Such Γ generalize the congruence subgroups of $\mathrm{SL}_2 \mathbb{Z}$ and are co-compact whenever B is ramified. For $\Gamma = \mathrm{SL}_2 \mathbb{Z}$, we prove that high energy bound eigenstates obey the Random Wave conjecture of Berry/Hejhal for third moments. In fact we show that the third moment of a wave's amplitude distribution decays as $E^{-\frac{1}{12}}$ in the high energy limit. In the more general case of maximal orders, we reduce an optimal quantitative version of the Quantum Unique Ergodicity conjecture of Rudnick–Sarnak to the Lindelöf Hypothesis, itself a consequence of the Riemann Hypothesis, for particular families of automorphic L -functions. Furthermore, our analysis shows that any lowering of the exponent in the Phragmén–Lindelöf convexity bound implies QUE. In the moment problem as well, a decisive role is played by ‘breaking convexity.’ That is to say, the boundaries of non-trivial exponents precisely agree when translated between physical and arithmetical formulations, for both of these problems.

We accomplish this translation by proving identities expressing triple-correlation integrals of eigenforms in terms of central values of the corresponding Rankin triple-product L -functions. Very general forms of such identities were proved by Harris–Kudla, and certain (more explicit) classical versions of these were given by Gross–Kudla and Böcherer–Schulze-Pillot, for definite quaternion algebras. In using the Harris–Kudla method to prove our own classical identities, we have to solve two main problems.

The first problem is to explicitly compute the adjoint of Shimizu's theta lift, which realizes the Jacquet–Langlands correspondence by transferring automorphic forms from GL_2 to $\mathrm{GO}(B)$, the latter being nearly the same as $B^\times \times B^\times$. As is well known, theta liftings from metaplectic to orthogonal groups are generally more difficult to characterize than lifts in the opposite direction, which can be evaluated directly in terms of Whittaker functions. Since $B^\times$ and hence $\mathrm{GO}(B)$ have ‘multiplicity-one’—as Jacquet–Langlands proved with the trace-formula—we are able to determine the adjoint of Shimizu's lift by duality, from explicit knowledge of Shimizu's lift itself. It is, however, necessary to generalize Shimizu's original calculations, since he only considered averages of lifts over isotypic bases of forms, which allowed him to employ Godement's theory of spherical functions. In order to deal with individual ramified forms, we replace this argument by explicit calculations involving Hecke operators. Thus we determine the Shimizu lifts of oldforms and newforms of square-free level, with (possibly imprimitive) neben-characters.

As a byproduct of these calculations, we obtain explicit formulas for all relevant GL_2 Whittaker functions. These play an important role in our second main problem: evaluation of Garrett/Piatetski-Shapiro–Rallis local zeta integrals in terms of the standard functorial triple-product L -factors. Our contribution lies in the calculation of archimedean zeta integrals with various types of ramification. In all of these ram-

ified cases, the canonical choices of local data for theta lifting are apparently useful only for evaluating *central* values of the associated zeta integrals. This was observed by Gross–Kudla in the non-archimedean case, and it appears to be a significant unexplained feature of the Harris–Kudla method. Finally, we re-prove (by elementary brute-force computation) the important result of Piatetski-Shapiro–Rallis on unramified non-archimedean zeta integrals, in anticipation of future generalizations to the many ramified non-archimedean cases.

Acknowledgements. This work is the product of my doctoral research at Princeton University, directed by Peter Sarnak. I am very grateful to him for explaining this fascinating problem to me, and for his generous help and encouragement, which cannot be overstated. I also thank Henryk Iwaniec and Goro Shimura for teaching me analytic and algebraic number theory, as well as Michael Boshernitzan and Yakov Sinai for teaching me ergodic theory. I gratefully acknowledge receiving funding through an NSF Graduate Fellowship and a Charlotte E. Proctor Honorific Fellowship. Finally, I thank my friends and family for their love and support.

Contents

Abstract	iii
Acknowledgements	iv
1 Automorphic Forms	1
1.1 Orders and Units of Quaternion Algebras	1
1.2 Automorphic Forms on Quaternion Groups	2
1.3 Hecke Operators	5
1.3.1 Double Cosets	5
1.3.2 Non-Archimedean Operators	8
1.3.3 Archimedean Operators	11
1.4 Automorphic Representations	12
1.5 Whittaker Models	16
2 Theta Lifting	21
2.1 Basic Setup	21
2.1.1 Groups	21
2.1.2 Measures	22
2.1.3 Dual-Pairs	24
2.2 Jacquet–Langlands Correspondence	25
2.2.1 Shimizu’s Theta Lift	25
2.2.2 Adjoint of Shimizu’s Lift	29
2.3 Local Data	31
3 L-functions	34
3.1 Local Langlands Correspondence	34
3.1.1 Adjoint Parameters	36
3.1.2 Triple Product Parameters	37
3.2 Zeta Integrals	39
3.2.1 Rankin–Selberg	39
3.2.2 Garrett/Piatetski-Shapiro–Rallis	42
4 Conclusion	51
4.1 Triple Product Identities	51
4.2 Applications to Quantum Chaos	52
4.2.1 Proofs	55

Chapter 1

Automorphic Forms

1.1 Orders and Units of Quaternion Algebras

Let B be a quaternion algebra over $\mathbb{Q}$. Then $B_v = B \otimes_{\mathbb{Q}} \mathbb{Q}_v$ is a quaternion algebra over $\mathbb{Q}_v$, for each place v of $\mathbb{Q}$. B is called ramified at v if B_v is a division algebra. We will always assume that B is unramified at ∞ , i.e. B is indefinite. Then the reduced discriminant d_B of B over $\mathbb{Q}$ equals the product of the finite even number of finite places p where B is ramified. The 2×2 matrix algebra $M = M_2$ over $\mathbb{Q}$ is totally unramified, and hence $d_M = 1$. We denote by D_v the unique division quaternion algebra over $\mathbb{Q}_v$, while M_v is the unique split (i.e. non-division) quaternion algebra.

We will frequently use the coordinate functions α_{ij} on $\alpha = \begin{bmatrix} \alpha_{11} & \alpha_{12} \\ \alpha_{21} & \alpha_{22} \end{bmatrix} \in M_v$, and also the notation $[X_{ij}] = \{\alpha \in M_v; \alpha_{ij} \in X_{ij} \subset \mathbb{Q}_v\}$.

Write the canonical anti-involution ι of B or B_v as $\alpha \mapsto \alpha^\iota$, noting its compatibility with field extensions, and define the reduced trace and norm by $\tau(\alpha) = \alpha + \alpha^\iota$ and $\nu(\alpha) = \alpha\alpha^\iota$. On M and M_v , ι is given explicitly as

$$\begin{bmatrix} a & b \\ c & d \end{bmatrix}^\iota = \begin{bmatrix} d & -b \\ -c & a \end{bmatrix},$$

so τ and ν correspond to the usual trace and determinant.

D_p has the valuation $\text{ord}_p \circ \nu$, and its valuation ring R_p is also the unique maximal order of D_p . Choose a uniformizer ϖ_p ; all bilateral R_p ideals in D_p are principal and of the form $\varpi_p^n R_p = R_p \varpi_p^n$, $n \in \mathbb{Z}$.

Every maximal order of M_p is conjugate to $M_2(\mathbb{Z}_p)$. We will be interested in those of the form $\begin{bmatrix} \mathbb{Z}_p & p^{-n}\mathbb{Z}_p \\ p^n\mathbb{Z}_p & \mathbb{Z}_p \end{bmatrix} = \begin{bmatrix} p^n & \\ & 1 \end{bmatrix}^{-1} M_2(\mathbb{Z}_p) \begin{bmatrix} p^n & \\ & 1 \end{bmatrix}$ for $n \in \mathbb{Z}$, although the set of all maximal orders is larger, having the structure of a $(p+1)$ -homogeneous tree for a natural notion of adjacency. It is a theorem of Hijikata that every Eichler order, an intersection of two maximal orders, is conjugate to $\begin{bmatrix} \mathbb{Z}_p & \mathbb{Z}_p \\ p^n\mathbb{Z}_p & \mathbb{Z}_p \end{bmatrix}$ for some $n \geq 0$. All Eichler orders containing the previous one are of the form $\begin{bmatrix} \mathbb{Z}_p & p^{-n}\mathbb{Z}_p \\ p^{\tilde{n}}\mathbb{Z}_p & \mathbb{Z}_p \end{bmatrix}$

for $0 \leq \dot{n} \leq \ddot{n} \leq n$.

Orders $\mathcal{O}$ of B exist and have the property that $\mathcal{O}_p := \mathcal{O} \otimes_{\mathbb{Z}} \mathbb{Z}_p$ is a maximal order in B_p for almost all p . Furthermore, for any other order $\mathcal{O}'$ of B , $\mathcal{O}'_p = \mathcal{O}_p$ for almost all p . Conversely, given a choice over all p of local orders $(\tilde{\mathcal{O}}_p)$ which satisfies $\tilde{\mathcal{O}}_p = \mathcal{O}_p$ for a.a. p , then $\mathcal{O}' := \{\alpha \in B; \alpha_p \in \tilde{\mathcal{O}}_p \forall p\}$ is an order in B with p -adic completions $\mathcal{O}'_p = \tilde{\mathcal{O}}_p$.

The adèle ring $B_{\mathbb{A}}$ is defined as the restricted direct product of all B_v relative to $\mathcal{O}_{\text{fin}} := \prod_p \mathcal{O}_p$, for any order $\mathcal{O}$ of B , and the idele group $B_{\mathbb{A}}^{\times}$ as the restricted direct product of all $B_v^{\times}$ relative to $\mathcal{O}_{\text{fin}}^{\times}$. B is contained diagonally in $B_{\mathbb{A}}$, and we will write $B = B_{\mathbb{Q}} \subset B_{\mathbb{A}}$ to emphasize this inculsion. By the previous paragraph, orders of B correspond 1-1 with compact open subgroups of B_{fin} .

Maximal orders in B are characterized as orders $\mathcal{O}$ s.t. every $\mathcal{O}_p$ is maximal. Choose a maximal order for each B , denoted $\mathcal{O}(d_B)$. Throughout the rest of the paper, for $p \nmid d_B$ we will identify B_p with M_p in such a way that $\mathcal{O}_p(d_B) = M_2(\mathbb{Z}_p)$. Furthermore, in the case $d_B = 1$ we identify B with M , and so $\mathcal{O}(1) = M_2(\mathbb{Z})$.

Now for $\dot{N}, \ddot{N} \in \mathbb{Q}^+$ relatively prime to d_B and s.t. $N = \dot{N}\ddot{N} \in \mathbb{Z}$, we define the order $\mathcal{O}(d_B, \vec{N})$ of B by

$$\mathcal{O}_p(d_B, \vec{N}) = \begin{cases} R_p & \text{if } p \mid d_B, \\ \begin{bmatrix} \mathbb{Z}_p & \dot{N}\mathbb{Z}_p \\ \ddot{N}\mathbb{Z}_p & \mathbb{Z}_p \end{bmatrix} & \text{if } p \nmid d_B. \end{cases}$$

Then $\mathcal{O}(d_B, \dot{N}, \dot{N}^{-1})$ is a maximal order, and

$$\mathcal{O}(d_B, \vec{N}) = \mathcal{O}(d_B, \dot{N}, \dot{N}^{-1}) \cap \mathcal{O}(d_B, \ddot{N}^{-1}, \ddot{N})$$

is an Eichler order. It has reduced discriminant $d_B N$ and is called of level N . Using strong approximation and Hijikata's result, it is easy to show that any Eichler order of level N in B is conjugate to $\mathcal{O}(d_B, (1, N))$. Note that the only Eichler orders containing $\mathcal{O}(d_B, \vec{N})$ are the $\mathcal{O}(d_B, \vec{N}')$ s.t. $\dot{N}' \mid \dot{N}$ and $\ddot{N}' \mid \ddot{N}$.

Throughout the rest of the paper, for any X possessing a character ν , we will use the notations

$$\begin{aligned} X_*^{(a)} &= \{\beta \in X_*; \nu(\beta) = a\}, \\ X_*^{(A)} &= \{\beta \in X_*; \nu(\beta) \in A\}, \\ X_{\infty}^{\pm} &= X_{\infty}^{(\mathbb{R}^{\pm})}, \quad X_p^{[a]} = X_p^{(a\mathbb{Z}_p^{\times})}. \end{aligned}$$

1.2 Automorphic Forms on Quaternion Groups

In this section, fix $\mathcal{O} = \mathcal{O}(d_B, \vec{N})$. Strong approximation for $\mathcal{O}_{\text{fin}}^{\times}$ is proved in [26]: $B_{\mathbb{A}}^{\times} = B_{\mathbb{Q}}^{\times} B_{\infty}^{+} \mathcal{O}_{\text{fin}}^{\times}$ since B is indefinite and $\nu(\mathcal{O}_{\text{fin}}^{\times}) = \mathbb{Z}_{\text{fin}}^{\times}$. Because of this, and since

$\mathcal{O}^{(1)} = B_{\mathbb{Q}}^{\times} \cap B_{\infty}^{+} \mathcal{O}_{\text{fin}}^{\times}$, there are homeomorphisms

$$\begin{aligned} B_{\mathbb{Q}}^{\times} \backslash B_{\mathbb{A}}^{\times} / \mathcal{O}_{\text{fin}}^{\times} &\simeq \mathcal{O}^{(1)} \backslash B_{\infty}^{+}, \\ B_{\mathbb{Q}}^{\times} \mathbb{R}^{+} \backslash B_{\mathbb{A}}^{\times} / \mathcal{O}_{\text{fin}}^{\times} &\simeq \mathcal{O}^{(1)} \backslash \text{PGL}_2^{+}(\mathbb{R}). \end{aligned}$$

Given a primitive Dirichlet character $\chi \bmod N^{\times}$, factor $\chi = \prod_p \chi_p$ and extend its domain to $\mathbb{Z}_{\text{fin}}^{\times}$ by means of the Chinese remainder theorem,

$$\begin{aligned} (\mathbb{Z}/N^{\times}\mathbb{Z})^{\times} &\simeq \prod_{p|N^{\times}} (\mathbb{Z}_p/N^{\times}\mathbb{Z}_p)^{\times} \\ &= \prod_{p|N^{\times}} \mathbb{Z}_p^{\times} / (1 + N^{\times}\mathbb{Z}_p). \end{aligned}$$

Using strong approximation, $\mathbb{A}^{\times} = \mathbb{Q}^{\times} \mathbb{R}^{+} \mathbb{Z}_{\text{fin}}^{\times}$, we define the grossen-character $\tilde{\chi}$ by $\tilde{\chi}(qrz) = \overline{\chi(z)}$. Note that $\tilde{\chi}_{\infty}(-1) = \chi(-1)$ and $\tilde{\chi}_p(p\mathbb{Z}_p^{\times}) = \chi(p)$ if $p \nmid N^{\times}$.

Since $\mathcal{O}$ has level N , the map $r_p : \mathcal{O}_p \rightarrow \mathbb{Z}_p/N\mathbb{Z}_p$ defined by $r_p(\beta) = [\beta_{11}]$ for $p \nmid d_B$, is a ring homomorphism, and hence so is $r : \mathcal{O}_{\text{fin}} \rightarrow \mathbb{Z}/N\mathbb{Z}$, $r(\beta) = (r_p(\beta_p))_{p|N}$. This permits us, for $N^{\times} \mid N$, to define $\tilde{\chi}$ on $\mathcal{O}_{\text{fin}}^{\times}$ and χ on $\mathcal{O}^{\times}$ by composition with r . (Note $\chi(\beta) = \overline{\tilde{\chi}(\beta)}$ for $\beta \in \mathcal{O}^{\times}$.) Since $\tilde{\chi}$ is well-defined on $\mathbb{Q}_{\text{fin}}^{\times} \cap \mathcal{O}_{\text{fin}}^{\times} = \mathbb{Z}_{\text{fin}}^{\times}$, it extends to a character of $\mathbb{Q}_{\text{fin}}^{\times} \mathcal{O}_{\text{fin}}^{\times}$.

As usual, $B_{\infty}^{\times} = \text{GL}_2(\mathbb{R})$ acts on $\mathbb{C} \backslash \mathbb{R}$ by linear fractional transformations,

$$z \mapsto \beta|z = \frac{az + b}{cz + d} \quad \text{for } \beta = \begin{bmatrix} a & b \\ c & d \end{bmatrix},$$

preserving the metric $ds^2 = (dx^2 + dy^2)/y^2$. Furthermore, B_{∞}^{+} acts on weight- k Maass forms ($k \in \mathbb{Z}$) on $\mathfrak{H} = \mathbb{R} + i\mathbb{R}^{+}$ by $\psi \mapsto \psi|_{\beta}^k$,

$$\psi|_{\beta}^k(z) = \left(\frac{cz+d}{cz+d} \right)^{-k} \psi(\beta|z).$$

Consider the space $A_k(d_B, \vec{N}, \chi)$ of smooth weight- k Maass forms ψ on $\mathfrak{H}$ which have moderate growth and satisfy the automorphy condition

$$\psi|_{\gamma_{\infty}}^k = \chi(\gamma) \psi \quad \text{for } \gamma \in \mathcal{O}^{(1)}(d_B, \vec{N}).$$

Since $-1 \in \mathcal{O}^{(1)}$, such ψ must vanish identically unless $\chi(-1) = (-1)^k$, so we assume this from now on. Let $C_k(d_B, \vec{N}, \chi) \subset A_k(d_B, \vec{N}, \chi)$ denote the cuspidal subspace, equipped with the Petersson inner-product

$$\langle \psi_1, \psi_2 \rangle := \int_{\mathcal{O}^{(1)} \backslash \mathfrak{H}} \psi_1(z) \overline{\psi_2(z)} \frac{dx dy}{y^2}.$$

In the adelic setting, we define $L^2(B_{\mathbb{Q}}^{\times} \backslash B_{\mathbb{A}}^{\times}, \tilde{\chi})$ as usual to consist of automorphic functions Ψ with central character $\tilde{\chi}$,

$$\Psi(\gamma z \beta) = \tilde{\chi}(z) \Psi(\beta) \quad \text{for } \gamma \in B_{\mathbb{Q}}^{\times}, z \in \mathbb{A}^{\times},$$

and finite norm under the inner-product

$$\langle \Psi_1, \Psi_2 \rangle := \frac{1}{2} \int_{PB_{\mathbb{Q}}^{\times} \backslash PB_{\mathbb{A}}^{\times}} \Psi_1(\beta) \overline{\Psi_2(\beta)} d^{\times} \beta.$$

The Tamagawa measure $d^{\times} \beta$ is defined in §2.1.2. Note that these two inner-products are normalized differently. We denote the right regular action of $\alpha \in B_{\mathbb{A}}^{\times}$ as

$$\alpha | \Psi(\beta) = \Psi(\beta \alpha) \quad \text{for } \Psi \in L^2(B_{\mathbb{Q}}^{\times} \backslash B_{\mathbb{A}}^{\times}, \tilde{\chi}),$$

and define $L_0^2(B_{\mathbb{Q}}^{\times} \backslash B_{\mathbb{A}}^{\times}, \tilde{\chi})$ as the closed invariant subspace of cuspidal Ψ .

Now let

$$K_v(d_B, \vec{N}) = \begin{cases} \mathcal{O}_p^{\times}(d_B, \vec{N}) & \text{if } v = p, \\ \text{SO}(2, \mathbb{R}) & \text{if } v = \infty, \end{cases}$$

and extend $\tilde{\chi}_{\infty}$ (dependent on k) to $\mathbb{R}^+ K_{\infty}$ from $\mathbb{R}^{\times}$ by setting

$$\tilde{\chi}_{\infty}(r \kappa_{\theta}) = e^{ik\theta} \quad \text{for } \kappa_{\theta} = \begin{bmatrix} \cos \theta & \sin \theta \\ -\sin \theta & \cos \theta \end{bmatrix}.$$

We define $\tilde{C}_k(d_B, \vec{N}, \tilde{\chi})$ to consist of all smooth $\Psi \in L_0^2(B_{\mathbb{Q}}^{\times} \backslash B_{\mathbb{A}}^{\times}, \tilde{\chi})$ satisfying

$$\kappa_v | \Psi = \tilde{\chi}_v(\kappa_v) \Psi \quad \text{for } \kappa_v \in K_v(d_B, \vec{N}).$$

(Note this is consistent with the inclusion of the center $\mathbb{A}^{\times} \subset B_{\mathbb{Q}}^{\times} \mathbb{R}^+ \mathcal{O}_{\text{fin}}^{\times}$.) Then we have an isomorphism $C_k(d_B, \vec{N}, \chi) \xrightarrow{\sim} \tilde{C}_k(d_B, \vec{N}, \tilde{\chi})$, given by

$$\Psi(\gamma \beta_{\infty}^+ \kappa_{\text{fin}}) = \tilde{\chi}(\kappa_{\text{fin}}) \psi|_{\beta_{\infty}^+}^k(\mathbf{i}).$$

If $\mathcal{O}(d_B, \vec{N}') \supset \mathcal{O}(d_B, \vec{N})$ and $N^{\times} \mid N'$, there is an inclusion

$$C_k(d_B, \vec{N}', \chi) \hookrightarrow C_k(d_B, \vec{N}, \chi).$$

Any form in the image of such a map with $pN' \mid N$ is called p -old, as are linear combinations of such forms, and any form orthogonal to all p -old forms is called p -new. Likewise at the archimedean place, we have raising and lowering operators,

$$R_k^{\pm} : C_k(d_B, \vec{N}, \chi) \longrightarrow C_{k \pm 2}(d_B, \vec{N}, \chi),$$

$$R_k^+ = (z - \bar{z}) \frac{\partial}{\partial z} + \frac{k}{2}, \quad R_k^- = (\bar{z} - z) \frac{\partial}{\partial \bar{z}} - \frac{k}{2}.$$

A weight- k form ψ is called ∞ -old if $\psi = R_{k \mp 2}^{\pm} \psi'$ for $\pm k \geq 0$, and ∞ -new if it is orthogonal to all ∞ -old forms. Since $R_{k \mp 2}^{\pm}$ and $R_k^{\mp}$ are adjoints, ψ is ∞ -new iff $R_k^{\mp} \psi = 0$ for $\pm k \geq 0$, and this is equivalent to $y^{-\frac{|k|}{2}} \psi(z)$ being a holomorphic/anti-holomorphic function of z . All forms with $k = 0$ are ∞ -old.

For the rest of this paper we assume that N is square-free, and factoring it into $N = N^b N^\sharp N^\chi$, we define $S_k(d_B, \vec{N}, \chi, N^b) \subset C_k(d_B, \vec{N}, \chi)$ as the subspace of forms which are p -old for $p \mid N^b$, p -new for $p \mid N^\sharp$, and ∞ -new if $|k| > 0$. We will also assume that $|k| \neq 1$.

1.3 Hecke Operators

In this section, fix $\mathcal{O} = \mathcal{O}(d_B, \vec{N})$ and χ primitive with conductor $N^\chi \mid N$, and assume that $N = N^b N^\sharp N^\chi$ is square-free. For $p \mid N^b N^\sharp$, set

$$\dot{\mathcal{O}}_p = \mathcal{O}_p(d_B, (\dot{N}, \frac{\ddot{N}}{p})), \quad \ddot{\mathcal{O}}_p = \mathcal{O}_p(d_B, (\frac{\dot{N}}{p}, \ddot{N})).$$

1.3.1 Double Cosets

For $\alpha \in B_p^\times$, define the double coset $T_p(\alpha) = \mathcal{O}_p^\times \alpha \mathcal{O}_p^\times$. If $p \mid d_B$, every $T_p(\alpha)$ is equal to some $T_p(\varpi_p^n) = \varpi_p^n R_p^\times = R_p^\times \varpi_p^n$ for $n \in \mathbb{Z}$, and $T_p(\varpi_p^n) \subset \mathcal{O}_p$ iff $n \geq 0$. Now for $p \nmid d_B$ and $\vec{a} \in (\mathbb{Q}_p^\times)^2$, define

$$T_p(\vec{a}) = T_p\left(\begin{bmatrix} \dot{a} & \\ & \ddot{a} \end{bmatrix}\right), \quad R_p(\vec{a}) = T_p\left(\begin{bmatrix} & \dot{a} \\ \ddot{a} & \end{bmatrix}\right).$$

Since all $a \in \mathbb{Q}_p^\times$ and $\alpha_{\vec{N}} = \begin{bmatrix} \dot{N} & \\ & \ddot{N} \end{bmatrix}$ normalize $\mathcal{O}_p^\times$,

$$T_p(a) = a \mathcal{O}_p^\times = \mathcal{O}_p^\times a, \quad R_p(\vec{N}) = \alpha_{\vec{N}} \mathcal{O}_p^\times = \mathcal{O}_p^\times \alpha_{\vec{N}}.$$

If $p \nmid d_B N$, every $T_p(\alpha)$ is equal to $T_p(p^{\vec{n}})$ for some $\dot{n} \geq \ddot{n}$ (set $n = \dot{n} - \ddot{n}$), and $T_p(p^{\vec{n}}) \subset \mathcal{O}_p$ iff $\ddot{n} \geq 0$. Given $\alpha \in B_p^\times$, the parameters $\dot{n}, \ddot{n}$ are determined by

$$\alpha \in p^{\ddot{n}}(\mathcal{O}_p \setminus p\mathcal{O}_p), \quad \nu(\alpha) \in p^{\dot{n}+\ddot{n}} \mathbb{Z}_p^\times.$$

Furthermore, we have the following left coset partition: (For right cosets, apply ι ; for lower triangular representatives, conjugate by $R_p(\vec{N})$.)

$$T_p(p^{\vec{n}}) = \bigcup_{m=0}^n \bigcup_{\substack{x \bmod p^m \\ p \nmid x \text{ if } 0 < m < n}} \left[\begin{array}{c} p^{m+\ddot{n}} \\ \dot{N} x p^{\ddot{n}} \\ p^{n-m+\ddot{n}} \end{array} \right] \mathcal{O}_p^\times.$$

We prove both well-known assertions of the previous paragraph at the same time.

Note that $T_p(p^{\vec{n}})$ contains the above union of left cosets, and they are disjoint since

$$\begin{aligned} & \begin{bmatrix} p^{m+\vec{n}} & \dot{N}xp^{\vec{n}} \\ & p^{n-m+\vec{n}} \end{bmatrix}^{-1} \begin{bmatrix} p^{m'+\vec{n}} & \dot{N}x'p^{\vec{n}} \\ & p^{n-m'+\vec{n}} \end{bmatrix} \\ &= \begin{bmatrix} p^{m'-m} & \dot{N}(x'p^{-m} - xp^{-m'}) \\ & p^{m-m'} \end{bmatrix} \notin \mathcal{O}_p^\times \end{aligned}$$

unless $m = m'$, $x \equiv x' \pmod{p^m}$. Now it suffices to show that the union of these left cosets over all $\dot{n} \geq \ddot{n} \geq 0$ has additive volume (see §2.1.2) equal to that of $\mathcal{O}_p \cap B_p^\times$, since for any $\alpha \in \mathcal{O}_p \cap B_p^\times$, $\text{vol}(\alpha\mathcal{O}_p^\times)/\text{vol}(\mathcal{O}_p^\times) = |\alpha|_{B_p} = |\nu(\alpha)|_p^2 > 0$. Finally,

$$\begin{aligned} \mathcal{O}_p \setminus \mathcal{O}_p^\times &= \begin{bmatrix} \mathbb{Z}_p^\times & \dot{N}\mathbb{Z}_p^\times \\ \dot{N}\mathbb{Z}_p^\times & \mathbb{Z}_p^\times \end{bmatrix}^{(p\mathbb{Z}_p)} \cup \begin{bmatrix} p\mathbb{Z}_p & \dot{N}p\mathbb{Z}_p \\ \dot{N}\mathbb{Z}_p^\times & \mathbb{Z}_p^\times \end{bmatrix} \cup \dots \cup \\ &\cup \begin{bmatrix} p\mathbb{Z}_p & \dot{N}p\mathbb{Z}_p \\ \dot{N}p\mathbb{Z}_p & \mathbb{Z}_p^\times \end{bmatrix} \cup \dots \cup \begin{bmatrix} p\mathbb{Z}_p & \dot{N}p\mathbb{Z}_p \\ \dot{N}p\mathbb{Z}_p & p\mathbb{Z}_p \end{bmatrix}, \quad \text{so} \end{aligned}$$

$$\begin{aligned} \text{vol}(\mathcal{O}_p^\times) &= 1 - \left(p^{-1} \left(\frac{p-1}{p} \right)^3 + 4p^{-2} \left(\frac{p-1}{p} \right)^2 + 4p^{-3} \left(\frac{p-1}{p} \right)^1 + p^{-4} \right) \\ &= (1 - p^{-1}) (1 - p^{-2}), \quad \text{and} \end{aligned}$$

$$\begin{aligned} \zeta_p(1)\zeta_p(2) &= \text{vol}(\mathcal{O}_p)/\text{vol}(\mathcal{O}_p^\times) \geq \text{vol}(\mathcal{O}_p \cap B_p^\times)/\text{vol}(\mathcal{O}_p^\times) \\ &\geq \sum_{\dot{n} \geq \ddot{n} \geq 0} (1 + (p-1)(1 + p + \dots + p^{\dot{n}-\ddot{n}-2}) + p^{\dot{n}-\ddot{n}}) (p^{-2\dot{n}-2\ddot{n}}) \\ &= \left(1 + (p-1) \sum_{n>0} p^{-n-1} \right) \sum_{\ddot{n} \geq 0} p^{-4\ddot{n}} \\ &= \left(\frac{1+p^{-2}}{1-p^{-1}} \right) \frac{1}{(1-p^{-4})} = \zeta_p(1)\zeta_p(2). \end{aligned}$$

If $p \mid N$, every $T_p(\alpha)$ (with $\alpha \in B_p^\times$) is equal to either $T_p(p^{\vec{n}})$ or $R_p(p^{\vec{n}}\vec{N})$ for some $\dot{n}, \ddot{n} \in \mathbb{Z}$, and $T_p(\alpha) \subset \mathcal{O}_p$ iff $\dot{n}, \ddot{n} \geq 0$. We distinguish among these as follows: First, $m = \min\{\dot{n}, \ddot{n}\}$ satisfies $\alpha \in p^m(\mathcal{O}_p \setminus p\mathcal{O}_p)$. Now determine the type of double coset and sign of $n = \dot{n} - \ddot{n}$ using the partition

$$\begin{aligned} \mathcal{O}_p \setminus p\mathcal{O}_p &= \begin{bmatrix} p\mathbb{Z}_p & \dot{N}\mathbb{Z}_p \\ \dot{N}\mathbb{Z}_p & \mathbb{Z}_p^\times \end{bmatrix}^{T,+} \cup \begin{bmatrix} \mathbb{Z}_p^\times & \dot{N}\mathbb{Z}_p \\ \dot{N}\mathbb{Z}_p & \mathbb{Z}_p^\times \end{bmatrix}^{T,0} \cup \begin{bmatrix} \mathbb{Z}_p^\times & \dot{N}\mathbb{Z}_p \\ \dot{N}\mathbb{Z}_p & p\mathbb{Z}_p \end{bmatrix}^{T,-} \cup \\ &\cup \begin{bmatrix} p\mathbb{Z}_p & \dot{N}p\mathbb{Z}_p \\ \dot{N}\mathbb{Z}_p^\times & p\mathbb{Z}_p \end{bmatrix}^{R,+} \cup \begin{bmatrix} p\mathbb{Z}_p & \dot{N}\mathbb{Z}_p^\times \\ \dot{N}\mathbb{Z}_p^\times & p\mathbb{Z}_p \end{bmatrix}^{R,0} \cup \begin{bmatrix} p\mathbb{Z}_p & \dot{N}\mathbb{Z}_p^\times \\ \dot{N}p\mathbb{Z}_p & p\mathbb{Z}_p \end{bmatrix}^{R,-}. \end{aligned}$$

The remaining parameter $\max\{\dot{n}, \ddot{n}\}$ is recovered from

$$\nu(\alpha) \in \begin{cases} p^{\dot{n}+\ddot{n}} \mathbb{Z}_p^\times & \text{if } \alpha \in T_p(p^{\ddot{n}}), \\ p^{1+\dot{n}+\ddot{n}} \mathbb{Z}_p^\times & \text{if } \alpha \in R_p(p^{\ddot{n}} \vec{N}). \end{cases}$$

These $T_p(\alpha)$, taking α diagonal or anti-diagonal, have left coset representatives as follows: (For right cosets, apply ι and conjugate by $R_p(\vec{N})$.)

$$T_p(\alpha) = \bigcup_{x \bmod p^{|\dot{n}|}} \alpha_x \mathcal{O}_p^\times, \quad \alpha_x = \begin{cases} \begin{bmatrix} 1 & \dot{N}x \\ & 1 \end{bmatrix} \alpha & \text{if } \dot{n} \geq 0, \\ \begin{bmatrix} 1 & \\ \ddot{N}x & 1 \end{bmatrix} \alpha & \text{if } \dot{n} \leq 0. \end{cases}$$

As in the case $p \nmid d_B N$, we prove all of these claims together by showing that the left cosets above (with $\dot{n}, \ddot{n} \geq 0$) completely partition $\mathcal{O}_p \cap B_p^\times$. The double cosets listed above are disjoint for unequal types/parameters by the preceding characterization, and the left cosets $\alpha_x \mathcal{O}_p^\times$ listed within the same double coset are disjoint, since $\alpha_x^{-1} \alpha_{x'} \notin \mathcal{O}_p^\times$ unless $x \equiv x' \bmod p^{|\dot{n}|}$.

Finally, $\mathcal{O}_p^\times = \begin{bmatrix} \mathbb{Z}_p^\times & \dot{N}\mathbb{Z}_p \\ \ddot{N}\mathbb{Z}_p & \mathbb{Z}_p^\times \end{bmatrix}$ has $\text{vol}(\mathcal{O}_p^\times) = p^{-1}(1 - p^{-1})^2$, and

$$\begin{aligned} \zeta_p(1)^2 &= \text{vol}(\mathcal{O}_p) / \text{vol}(\mathcal{O}_p^\times) \geq \text{vol}(\mathcal{O}_p \cap B_p^\times) / \text{vol}(\mathcal{O}_p^\times) \\ &\geq \sum_{\ddot{n} \geq 0} p^{-4\ddot{n}} + 2 \sum_{\dot{n} > \ddot{n} \geq 0} p^{\dot{n}-\ddot{n}-2\dot{n}-2\ddot{n}} \\ &\quad + \sum_{\ddot{n} \geq 0} p^{-4\ddot{n}-2} + 2 \sum_{\dot{n} > \ddot{n} \geq 0} p^{\dot{n}-\ddot{n}-2\dot{n}-2\ddot{n}-2} \\ &= (1 + p^{-2}) \left(1 + 2 \sum_{n > 0} p^{-n} \right) \sum_{\ddot{n} \geq 0} p^{-4\ddot{n}} \\ &= (1 + p^{-2}) \left(\frac{1 + p^{-1}}{1 - p^{-1}} \right) \frac{1}{(1 - p^{-4})} = \zeta_p(1)^2. \end{aligned}$$

In the subcase $p \mid N^\flat N^\sharp$, we write double cosets of $\dot{\mathcal{O}}_p^\times, \ddot{\mathcal{O}}_p^\times$ as $\dot{T}_p(\alpha), \ddot{T}_p(\alpha)$. It is easy to check by the disjointness/volume argument that

$$\begin{aligned} \dot{T}_p(1) &= T_p(1) \cup R_p(\dot{N}, \frac{\ddot{N}}{p}), \\ \ddot{T}_p(1) &= T_p(1) \cup R_p(\frac{\dot{N}}{p}, \ddot{N}). \end{aligned}$$

1.3.2 Non-Archimedean Operators

The space $\mathcal{S}(B_p^\times)$ of $\mathbb{C}$ -valued Schwarz functions on $B_p^\times$ forms an algebra under convolution (using the measure $d_p^\times\beta$ defined in §2.1.2),

$$(\phi_p * \phi'_p)(\alpha) := \int_{B_p^\times} \phi_p(\alpha\beta^{-1}) \phi'_p(\beta) d_p^\times\beta,$$

and it acts on $L_0^2(B_\mathbb{Q}^\times \backslash B_\mathbb{A}^\times, \tilde{\chi})$ via the right regular representation,

$$\phi_p | \Psi := \int_{B_p^\times} \beta | \Psi \phi_p(\beta) d_p^\times\beta.$$

Now we will define and study the subquotient algebras $\mathcal{H}_p^* = \mathcal{H}_p, \mathcal{H}_p^\flat, \mathcal{H}_p^\sharp, \mathcal{H}_p^\chi$, in the respective cases $p \nmid N, p \mid N^\flat, p \mid N^\sharp, p \mid N^\chi$, acting on $\tilde{S}_k(d_B, \vec{N}, \tilde{\chi}, N^\flat)$.

For any p , define $\mathcal{H}_p$ to be the subalgebra of $\mathcal{S}(B_p^\times)$ generated by the functions

$$\phi_{T_p(\alpha)} := \frac{1}{\text{vol}^\times(\mathcal{O}_p^\times)} \mathbb{I}_{T_p(\alpha)} \quad \text{with } \alpha \in B_p^\times.$$

We abuse notation by writing $T_p(\alpha)$ in place of $\phi_{T_p(\alpha)}$. For $a \in \mathbb{Q}_p^\times$, let

$$T_p^{[a]} := \sum_{\alpha \in \mathcal{O}_p^\times \backslash \mathcal{O}_p^{[a]} / \mathcal{O}_p^\times} T_p(\alpha) \in \mathcal{H}_p.$$

If $p \nmid N^\chi$, $\mathcal{H}_p$ acts on $\tilde{S}_k(d_B, \vec{N}, \tilde{\chi}, N^\flat)$ non-trivially. The element

$$T_p(\alpha)^\vee := T_p(\alpha^{-1}) \in \mathcal{H}_p$$

acts as the adjoint of $T_p(\alpha)$, and $T_p(p)$ acts as the scalar $\chi(p)$.

If $p \mid d_B$ then $\mathcal{H}_p \simeq \mathbb{C}(T_p(\varpi_p))$ and $T_p^{[p^n]} = \begin{cases} T_p(\varpi_p)^n & \text{if } n \geq 0, \\ 0 & \text{if } n < 0. \end{cases}$

If $p \nmid d_B N$, it is well known (see [36]) that $\mathcal{H}_p \simeq \mathbb{C}[T_p(p, 1), T_p(p), T_p(p)^{-1}]$,

$$T_p^{[p^n]} = \sum_{\substack{\vec{n} + \vec{n} = n \\ \vec{n} \geq \vec{0}}} T_p(p^{\vec{n}}),$$

$$\mathcal{H}_p[[X]] \ni \sum_{n=0}^{\infty} T_p^{[p^n]} X^n = (T_p(1) - T_p(p, 1)X + p T_p(p)X^2)^{-1}.$$

For $p \mid N$, we begin by computing the multiplication rules in $\mathcal{H}_p$:

$$T_p(\vec{a})^n = T_p(\vec{a}^n) \text{ for } n \geq 0, \quad R_p(\vec{N})^2 = T_p(p),$$

$$T_p(\vec{a}) T_p(p) = T_p(\vec{a}p) = T_p(p) T_p(\vec{a}),$$

$$T_p(\vec{a}) R_p(\vec{N}) = R_p(\vec{a}\vec{N}) = R_p(\vec{N}) T_p(\vec{a}, \vec{a}),$$

$$\begin{aligned} T_p(p, 1) T_p(1, p) &= p T_p(p) + (p-1) T_p(p, 1) R_p(\vec{N}), \\ T_p(1, p) T_p(p, 1) &= p T_p(p) + (p-1) T_p(1, p) R_p(\vec{N}). \end{aligned}$$

Also,

$$T_p^{[p^n]} = \sum_{\substack{\dot{n} + \ddot{n} = n \\ \dot{n}, \ddot{n} \geq 0}} T_p(p^{\vec{n}}) + \sum_{\substack{1 + \dot{n} + \ddot{n} = n \\ \dot{n}, \ddot{n} \geq 0}} R_p(p^{\vec{n}} \vec{N}).$$

Now define $\mathcal{H}_p^\flat$ as the centralizer of $R_p(\vec{N})$ in $\mathcal{H}_p$. Shimizu proved in [35] (see also [28]) that $\mathcal{H}_p^\flat$ is a maximal commutative subalgebra of $\mathcal{H}_p$, that it contains all $T_p^{[p^n]}$, and that

$$\sum_{n=0}^{\infty} T_p^{[p^n]} X^n = \frac{T_p(1) + p R_p(\vec{N}) X}{T_p(1) - T_p^{[p]} X + p R_p(\vec{N}) X + p T_p(p) X^2}.$$

To define $\mathcal{H}_p^\sharp$, we first note that

$$\begin{aligned} \dot{T}_p(1) &= \frac{1}{p+1} (T_p(1) + R_p(\dot{N}, \frac{\ddot{N}}{p})), \\ \ddot{T}_p(1) &= \frac{1}{p+1} (T_p(1) + R_p(\frac{\dot{N}}{p}, \ddot{N})), \end{aligned}$$

and compute the relations

$$\begin{aligned} T_p(1, p) \dot{T}_p(1) &= \frac{p}{p+1} (T_p(1, p) + R_p(\vec{N})) = \dot{T}_p(1) T_p(1, p), \\ T_p(p, 1) \ddot{T}_p(1) &= \frac{p}{p+1} (T_p(p, 1) + R_p(\vec{N})) = \ddot{T}_p(1) T_p(p, 1). \end{aligned}$$

If $p \mid N^\sharp$, the elements $\dot{T}_p(1)$, $\ddot{T}_p(1)$ of $\mathcal{H}_p$ annihilate $\tilde{S}_k(d_B, \vec{N}, \tilde{\chi}, N^\flat)$, while $T_p(1, p)$, $T_p(p, 1)$ are invertible, as we will see in §1.4. Such a representation of $\mathcal{H}_p$ factors through the quotient

$$\begin{aligned} \mathcal{H}_p^\sharp &:= \mathcal{H}_p / (T_p(1, p) + R_p(\vec{N})) \\ &= \mathcal{H}_p / (T_p(p, 1) + R_p(\vec{N})) \simeq \mathbb{C}(R_p^\sharp(\vec{N})), \\ T_p^{[p^n]} &\equiv \begin{cases} (-R_p(\vec{N}))^n & \text{if } n \geq 0, \\ 0 & \text{if } n < 0, \end{cases} \quad \text{in } \mathcal{H}_p^\sharp. \end{aligned}$$

Now suppose $p \mid N^\chi$. For $\beta \in B_p^\times$, $\delta = \det \beta$, $m \in \{1, 2\}$, $m' = 3 - m$, we define

$$\begin{aligned} \phi_{T_p(p^{(n_1, n_2)})}^{\chi m, m}(\beta) &= \frac{1}{\text{vol}^\times(\mathcal{O}_p^\times)} \mathbb{I}_{T_p(p^{(n_1, n_2)})}(\beta) \begin{cases} \overline{\tilde{\chi}_p(\beta_{mm})} & \text{if } n_m \leq n_{m'}, \\ \tilde{\chi}_p(\delta / \beta_{m'm'}) & \text{if } n_m \geq n_{m'}, \end{cases} \\ \phi_{T_p(p^{(n_1, n_2)})}^{\chi m, m'}(\beta) &= 0, \end{aligned}$$

$$\begin{aligned}\phi_{R_p(p^{(n_1, n_2)} \vec{N})}^{\chi m, m'}(\beta) &= \frac{1}{\text{vol}^\times(\mathcal{O}_p^\times)} \mathbb{I}_{R_p(p^{(n_1, n_2)} \vec{N})}(\beta) \begin{cases} \overline{\tilde{\chi}_p(\beta_{mm'})} & \text{if } n_m \leq n_{m'}, \\ \tilde{\chi}_p(-\delta/\beta_{m'm}) & \text{if } n_m \geq n_{m'}, \end{cases} \\ \phi_{R_p(p^{(n_1, n_2)} \vec{N})}^{\chi m, m}(\beta) &= 0.\end{aligned}$$

These cases were treated somewhat differently by Miyake in [26, 27]. It is straightforward to check, using the partition in §1.3.1, that

$$\phi_{T_p(\alpha)}^{\chi \vec{m}}(\dot{\kappa} \beta \ddot{\kappa}) = \overline{\tilde{\chi}_p(\dot{\kappa} \dot{m} \ddot{\kappa} \ddot{m})} \phi_{T_p(\alpha)}^{\chi \vec{m}}(\beta) \quad \text{for } \dot{\kappa}, \ddot{\kappa} \in \mathcal{O}_p^\times.$$

Any function $\phi^\chi \in \mathcal{S}(B_p^\times)$ satisfying this transformation property and supported on a double-coset $T_p(\alpha) \in \mathcal{H}_p$ is determined by its value $\phi^\chi(\beta)$ on any $\beta \in T_p(\alpha)$. By taking β to be diagonal or anti-diagonal and considering the left and right actions of diagonal $\kappa \in \mathcal{O}_p^\times$, we see that in the diagonal case, $\dot{m} \neq \ddot{m} \Rightarrow \phi^\chi = 0$, and in the anti-diagonal case, $\dot{m} = \ddot{m} \Rightarrow \phi^\chi = 0$. Thus, any such ϕ^χ is proportional to the corresponding $\phi_{T_p(\alpha)}^{\chi \vec{m}}$ listed above. These functions are normalized so that their values are multiplicative,

$$\phi_{T_p(\alpha_1)}^{\chi \vec{m}_1}(\alpha_1) \cdot \phi_{T_p(\alpha_2)}^{\chi \vec{m}_2}(\alpha_2) = \phi_{T_p(\alpha_1 \alpha_2)}^{\chi \vec{m}_1, \vec{m}_2}(\alpha_1 \alpha_2),$$

for diagonal or anti-diagonal $\alpha_j \in B_p^\times$, whenever neither side vanishes. Now it follows from convolving supports in $\mathcal{H}_p$ that the ϕ^χ convolve as

$$\begin{aligned}(\phi_{T_p(\vec{a})}^{\chi m, m})^{*n} &= \phi_{T_p(\vec{a}^n)}^{\chi m, m} \quad \text{for } n \geq 0, & \phi_{R_p(\vec{N})}^{\chi m, m'} * \phi_{R_p(\vec{N})}^{\chi m', m} &= \phi_{T_p(p)}^{\chi m, m}, \\ \phi_{T_p(\vec{a})}^{\chi m, m} * \phi_{T_p(p)}^{\chi m, m} &= \phi_{T_p(\vec{a}p)}^{\chi m, m} = \phi_{T_p(p)}^{\chi m, m} * \phi_{T_p(\vec{a})}^{\chi m, m}, \\ \phi_{T_p(\vec{a})}^{\chi m, m} * \phi_{R_p(\vec{N})}^{\chi m, m'} &= \phi_{R_p(\vec{a}\vec{N})}^{\chi m, m'} = \phi_{R_p(\vec{N})}^{\chi m, m'} * \phi_{T_p(\vec{a}, \vec{a})}^{\chi m', m'}, \\ \phi_{T_p(p, 1)}^{\chi m, m} * \phi_{T_p(1, p)}^{\chi m, m} &= p \phi_{T_p(p)}^{\chi m, m} = \phi_{T_p(1, p)}^{\chi m, m} * \phi_{T_p(p, 1)}^{\chi m, m}.\end{aligned}$$

Define $\mathcal{H}_p^\chi$ to be the $\mathbb{C}$ -algebra of finite linear combinations of the functions

$$T_p^\chi(\vec{a}) := \phi_{T_p(\vec{a})}^{\chi 1, 1} + \phi_{T_p(\vec{a})}^{\chi 2, 2} \quad \text{for } \vec{a} \in (\mathbb{Q}_p^\times)^2,$$

$$\text{so } \mathcal{H}_p^\chi \simeq \mathbb{C}(T_p^\chi(p, 1), T_p^\chi(1, p)).$$

Also define

$$R_p^\chi(\vec{a}\vec{N}) = \phi_{R_p(\vec{a}\vec{N})}^{\chi 1, 2} + \phi_{R_p(\vec{a}\vec{N})}^{\chi 2, 1}.$$

All of the functions ϕ^χ satisfy $\phi^\chi(z\beta) = \overline{\tilde{\chi}_p(z)} \phi^\chi(\beta)$ for $z \in \mathbb{Z}_p^\times$, and hence act non-trivially on $L_0^2(B_\mathbb{Q}^\times \backslash B_\mathbb{A}^\times, \tilde{\chi})$.

Given $\eta^\chi \in \prod_{p|N^\chi}(\mathbb{Z}/2\mathbb{Z})$, define

$$\begin{aligned}\Psi^{\eta^\chi} &= \left(\prod_{p|N^\chi} R_p^\chi(\vec{N})^{\eta_p^\chi} \right) \Psi \quad \text{and} \\ \tilde{C}_k^{\eta^\chi}(d_B, \vec{N}, \tilde{\chi}) &= \left(\prod_{p|N^\chi} R_p^\chi(\vec{N})^{\eta_p^\chi} \right) \tilde{C}_k(d_B, \vec{N}, \tilde{\chi}),\end{aligned}$$

which consists of $\Psi \in L_0^2(B_\mathbb{Q}^\times \backslash B_\mathbb{A}^\times, \tilde{\chi})$ satisfying

$$\kappa|\Psi = \begin{cases} \tilde{\chi}_p(\kappa_{11}) \Psi & \text{if } \eta_p^\chi = 0, \\ \tilde{\chi}_p(\kappa_{22}) \Psi & \text{if } \eta_p^\chi = 1, \end{cases} \quad \text{for } \kappa \in \mathcal{O}_p^\times, p \mid N^\chi,$$

plus the usual K_v condition for $v \nmid N^\chi$. The action of $T_p^\chi(\vec{a}) \in \mathcal{H}_p^\chi$ stabilizes each $\tilde{C}_k^{\eta^\chi}(d_B, \vec{N}, \tilde{\chi})$ and is adjoint to the action of $T_p^\chi(\vec{a})^\vee := T_p^\chi(\vec{a}^{-1})$, while $T_p^\chi(p)$ acts as the identity.

1.3.3 Archimedean Operators

Let $\mathcal{H}_\infty^\star$ denote the semigroup generated by T_∞^Δ alone,

$$\begin{aligned}T_\infty^\Delta \Psi &= \left(-y^2 \left(\frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2} \right) - y \frac{\partial^2}{\partial x \partial \theta} \right) \sigma_z \kappa_\theta |\Psi, \\ T_\infty^- \Psi &= \epsilon_\infty |\Psi, \quad \epsilon_\infty = \begin{bmatrix} 1 & \\ & -1 \end{bmatrix} \in B_\infty^\times.\end{aligned}$$

It is easy to check that these commute and

$$\begin{aligned}T_\infty^\Delta : \tilde{C}_k^{\eta^\chi}(d_B, \vec{N}, \tilde{\chi}) &\rightarrow \tilde{C}_k^{\eta^\chi}(d_B, \vec{N}, \tilde{\chi}), \\ T_\infty^- : \tilde{C}_k^{\eta^\chi}(d_B, \vec{N}, \tilde{\chi}) &\rightarrow \tilde{C}_{-k}^{\eta^\chi}(d_B, \vec{N}, \tilde{\chi}).\end{aligned}$$

For $\eta \in \prod_v(\mathbb{Z}/2\mathbb{Z})$, $\eta^\chi = (\eta_p)_{p|N^\chi}$, define

$$\begin{aligned}\Psi^\eta &= (T_\infty^-)^{\eta_\infty} \Psi^{\eta^\chi}, \\ \tilde{C}_k^\eta(d_B, \vec{N}, \tilde{\chi}) &= \tilde{C}_{(-1)^{\eta_\infty} k}^{\eta^\chi}(d_B, \vec{N}, \tilde{\chi}).\end{aligned}$$

Now consider the $\mathbb{C}$ -valued coordinates,

$$\begin{aligned}X(\beta) &= \frac{1}{2}(a+d) + \frac{i}{2}(b-c), \\ Y(\beta) &= \frac{1}{2}(a-d) + \frac{i}{2}(b+c),\end{aligned} \quad \beta = \begin{bmatrix} a & b \\ c & d \end{bmatrix} \in B_\infty,$$

in which we write the $(2,2)$ -form ν and a positive-definite majorant P ,

$$\begin{aligned}\nu(\beta) &= \frac{1}{2} \text{tr}(\beta \beta^t) = X\overline{X} - Y\overline{Y} = ad - bc, \\ P(\beta) &:= \frac{1}{2} \text{tr}(\beta \beta^t) = X\overline{X} + Y\overline{Y} = \frac{1}{2}(a^2 + b^2 + c^2 + d^2).\end{aligned}$$

It is straightforward to compute, writing $B_\infty^+ \ni \beta = \kappa_{\dot{\theta}} \begin{bmatrix} re^{t/2} & \\ & re^{-t/2} \end{bmatrix} \kappa_{\ddot{\theta}}$, that

$$\begin{aligned} X(\beta) &= e^{i(+\dot{\theta}+\ddot{\theta})} r \cosh \frac{t}{2}, & \nu(\beta) &= r^2, \\ Y(\beta) &= e^{i(-\dot{\theta}+\ddot{\theta})} r \sinh \frac{t}{2}, & P(\beta) &= r^2 \cosh t. \end{aligned}$$

Now define $\phi_{X_\infty}^{k,\xi}, \phi_{Y_\infty}^{k,\xi} \in \mathcal{S}(\mathbb{R}^+ \setminus B_\infty^+)$ by

$$\begin{aligned} \phi_{X_\infty}^{k,\xi}(\beta) &= \frac{1}{\pi} \left(\overline{X}(\beta) / \nu(\beta)^{\frac{1}{2}} \right)^{\frac{k}{2}} e(i\xi P(\beta) / \nu(\beta)), \\ &= \frac{1}{\pi} \overline{\tilde{\chi}(\kappa_{\dot{\theta}}) \tilde{\chi}(\kappa_{\ddot{\theta}})} (\cosh \frac{t}{2})^{|k|} e(i\xi \cosh t), \\ \phi_{Y_\infty}^{k,\xi}(\beta) &= \frac{1}{\pi} \left(\overline{Y}(\beta) / \nu(\beta)^{\frac{1}{2}} \right)^{\frac{k}{2}} e(i\xi P(\beta) / \nu(\beta)), \\ &= \frac{1}{\pi} \tilde{\chi}(\kappa_{\dot{\theta}}) \overline{\tilde{\chi}(\kappa_{\ddot{\theta}})} (\sinh \frac{t}{2})^{|k|} e(i\xi \cosh t), \end{aligned} \quad z^{\frac{k}{2}} = \begin{cases} z^k & \text{if } k > 0, \\ 1 & \text{if } k = 0, \\ \bar{z}^{|k|} & \text{if } k < 0. \end{cases}$$

These give rise to Hecke operators again commuting with T_∞^Δ ,

$$\begin{aligned} T_{*\infty}^{k,\xi} : C_k^{\eta^\times}(d_B, \vec{N}, \tilde{\chi}) &\rightarrow C_k^\eta(d_B, \vec{N}, \tilde{\chi}), \\ T_{*\infty}^{k,\xi} \Psi &= \int_{\mathbb{R}^+ \setminus B_\infty^+} \beta |\Psi| \phi_{*\infty}^{k,\xi}(\beta) d_\infty^\times \beta, \end{aligned} \quad \eta_\infty = \begin{cases} 0 & \text{if } * = X, \\ 1 & \text{if } * = Y. \end{cases}$$

1.4 Automorphic Representations

Given $\Psi \in \tilde{S}_k(d_B, \vec{N}, \tilde{\chi}, N^b)$, define $\pi \subset L_0^2(B_\mathbb{Q}^\times \setminus B_\mathbb{A}^\times, \tilde{\chi})$ as the subspace of $K_\mathbb{A}(d_B)$ -finite vectors in the closure of the span of $\{\beta |\Psi|; \beta \in B_\mathbb{A}^\times\}$. This is the cuspidal automorphic representation of $B_\mathbb{A}^\times$ attached to Ψ . If π is irreducible, then $\pi = \tilde{\otimes}_v \pi_v$ and $\Psi = \tilde{\otimes}_v \Psi_v$ factor as restricted tensor products over all places v , relative to a choice of spherical unit vectors Ψ_v at almost all places. In this section, we recall the classification of pre-unitary irreducible admissible representations of $B_v^\times$ having square-free conductor [17, 6, 4], and explicitly compute the actions of Hecke operators on these models. As a consequence, we have

Lemma 1. *The eigenforms in $\tilde{S}_k(d_B, \vec{N}, \tilde{\chi}, N^b)$ of $\mathcal{H}^\star$ comprise an orthogonal basis, and their tuples of eigenvalues have multiplicity one. The corresponding automorphic representations are each shared by $2^{\#\{p|N^b\}}$ basis elements, and these have the same eigenvalues for all $v \nmid N^b$.*

Proof This follows from multiplicity-one for automorphic representations of $B_\mathbb{A}^\times$ (proved by Jacquet–Langlands using the trace-formula) and Casselman’s theorem, plus the local calculations below for $p \mid N^b$. $\square$

For $p \nmid d_B N$, π_p is equivalent to the unramified continuous series $\pi(|\cdot|_p^{\frac{s_p}{2}}, |\cdot|_p^{\frac{\bar{s}_p}{2}})$ with

central character $\tilde{\chi}_p$. Since π_p is pre-unitary, either

$$\begin{aligned}\dot{s}_p &= -\ddot{s}_p - \log_p(\chi(p)) \in i\mathbb{R}/\frac{2\pi i}{\log(p)}\mathbb{Z}, \quad \text{or} \\ \dot{s}_p &= -\overline{\ddot{s}_p} = \sigma_p + it_p, \quad \text{with} \\ \sigma_p &\in (0, \frac{1}{2}), \quad t_p \in \frac{i}{2} \log_p(\chi(p)) + \frac{\pi}{\log(p)}\mathbb{Z}/\frac{2\pi}{\log(p)}\mathbb{Z}.\end{aligned}$$

Casselman's theorem implies that the right $\mathcal{O}_p^\times$ -invariant vector Ψ_p is unique up to scaling in π_p , and hence corresponds to $V_p^0 \in \pi(|\dot{s}_p|, |\ddot{s}_p|)$ of the form

$$V_p^0 \left(\begin{bmatrix} \dot{a} & * \\ & \ddot{a} \end{bmatrix} \kappa \right) = |\dot{a}|_p^{\dot{s}_p} |\ddot{a}|_p^{\ddot{s}_p} \left| \frac{\dot{a}}{\ddot{a}} \right|_p^{\frac{1}{2}} V_p^0(1) \quad \text{for } \kappa \in \mathcal{O}_p^\times.$$

In terms of the parameters, Ψ_p has $T_p(p, 1)$ eigenvalue $\lambda_p = p^{\frac{1}{2}} (p^{-\dot{s}_p} + p^{-\ddot{s}_p})$ and $T_p(p)$ eigenvalue $\chi(p) = p^{-\dot{s}_p - \ddot{s}_p}$, and hence by the analysis in §1.3.2, $T_p^{[p^n]}$ eigenvalue

$$\begin{aligned}\lambda_{p^n} &= p^{\frac{n}{2}} (p^{-n\dot{s}_p} + p^{-(n-1)\dot{s}_p - \ddot{s}_p} + \dots + p^{-n\ddot{s}_p}) \\ &= p^{\frac{n}{2}} \left(\frac{p^{-(n+1)\dot{s}_p} - p^{-(n+1)\ddot{s}_p}}{p^{-\dot{s}_p} - p^{-\ddot{s}_p}} \right).\end{aligned}$$

If $p \mid N^\flat$, π_p is again equivalent to an unramified continuous series, exactly as above. However by Casselman's theorem, the subspace of right $\mathcal{O}_p^\times$ -invariant vectors is spanned by a right $\dot{\mathcal{O}}_p^\times$ -invariant vector $\dot{V}_p^0$ and a right $\ddot{\mathcal{O}}_p^\times$ -invariant vector $\ddot{V}_p^0$, each unique up to scaling. We relate their normalizations by defining

$$\begin{aligned}\ddot{V}_p^0 &= \begin{bmatrix} 1 & \\ & p \end{bmatrix} \dot{V}_p^0 = R_p(\vec{N}) \dot{V}_p^0, \\ \implies \chi(p) \dot{V}_p^0 &= T_p(p) \dot{V}_p^0 = R_p(\vec{N}) \ddot{V}_p^0.\end{aligned}$$

Using coset representatives, we see that

$$\begin{aligned}T_p(p, 1) \dot{V}_p^0 &= \lambda_p \dot{V}_p^0 - \ddot{V}_p^0, \\ T_p(p, 1) \ddot{V}_p^0 &= p \chi(p) \dot{V}_p^0.\end{aligned}$$

We have now determined the actions of $R_p(\vec{N})$ and $T_p(p, 1)$, and hence also

$$\begin{aligned}T_p(1, p) &= R_p(\vec{N})^{-1} T_p(p, 1) R_p(\vec{N}), \\ T_p^{[p]} &= T_p(p, 1) + T_p(1, p) + R_p(\vec{N}).\end{aligned}$$

With respect to the basis $[\dot{V}_p^0, \ddot{V}_p^0]$, these operators are represented by the matrices

$$\begin{aligned} T_p(p, 1) &\sim \begin{bmatrix} \lambda_p & p\chi(p) \\ -1 & \end{bmatrix}, & T_p(1, p) &\sim \begin{bmatrix} & -\chi(p) \\ p & \lambda_p \end{bmatrix}, \\ T_p^{[p]} &\sim \begin{bmatrix} \lambda_p & p\chi(p) \\ p & \lambda_p \end{bmatrix}, & R_p(\vec{N}) &\sim \begin{bmatrix} & \chi(p) \\ 1 & \end{bmatrix}. \end{aligned}$$

The latter two are normal and commute, and so have orthogonal eigenvectors

$$\begin{aligned} V_p^b &:= \dot{V}_p^0 + \overline{\varepsilon_p} \ddot{V}_p^0, & \varepsilon_p^2 &= \chi(p), \\ T_p^{[p]} V_p^b &= (\lambda_p + p\varepsilon_p) V_p^b, & R_p(\vec{N}) V_p^b &= \varepsilon_p V_p^b. \end{aligned}$$

Then by the considerations of §1.3.2,

$$T_p^{[p^n]} V_p^b = (\lambda_{p^n} + p\varepsilon_p \lambda_{p^{n-1}}) V_p^b.$$

The idempotents $\dot{T}_p(1)$, $\ddot{T}_p(1)$ act as orthogonal projectors to $\mathbb{C} \dot{V}_p^0$, $\mathbb{C} \ddot{V}_p^0$, and using the above calculations we see that $\dot{T}_p(1) \ddot{V}_p^0 = \frac{\lambda_p}{p+1} \dot{V}_p^0$, so

$$\langle \ddot{V}_p^0, \dot{V}_p^0 \rangle = \frac{\lambda_p}{p+1} = \chi(p) \langle \dot{V}_p^0, \ddot{V}_p^0 \rangle, \quad \langle V_p^b, V_p^b \rangle = 2(1 + \frac{\overline{\varepsilon_p} \lambda_p}{p+1}).$$

For $p \mid N^\sharp$, π_p is equivalent to the special representation $\sigma(|\dot{s}_p|, |\ddot{s}_p|)$, where

$$\dot{s}_p = -\overline{\ddot{s}_p} = \frac{1}{2} + it_p, \quad t_p \in \frac{i}{2} \log_p(\chi(p)) + \frac{\pi}{\log(p)} \mathbb{Z} / \frac{2\pi}{\log(p)} \mathbb{Z}.$$

This is an irreducible invariant subspace of $\text{Ind}_{T_p}^{G_p}(|\dot{s}_p|, |\ddot{s}_p|)$, consisting of V_p s.t.

$$\int_{\mathbb{Q}_p} V_p \left(\begin{bmatrix} 1 & \\ x & 1 \end{bmatrix} \right) dx = 0.$$

For an $\mathcal{O}_p^\times$ -invariant vector $V_p^\sharp$, this is equivalent to

$$|\ddot{N}|_p V_p^\sharp(1) + |\dot{N}|_p V_p^\sharp \left(\begin{bmatrix} & 1 \\ 1 & \end{bmatrix} \right) = 0,$$

and so for $\kappa \in \mathcal{O}_p^\times$,

$$\begin{aligned} V_p^\sharp \left(\begin{bmatrix} \dot{a} & * \\ & \ddot{a} \end{bmatrix} \kappa \right) &= |\dot{a}\ddot{a}|_p^{it_p} \left| \frac{\dot{a}}{\ddot{a}} \right|_p V_p^\sharp(1), \\ V_p^\sharp \left(\begin{bmatrix} * & \dot{a} \\ & \ddot{a} \end{bmatrix} \kappa \right) &= -|\dot{a}\ddot{a}|_p^{it_p} \left| \frac{\dot{a}\ddot{N}}{\ddot{a}\dot{N}} \right|_p V_p^\sharp(1). \end{aligned}$$

Casselman's theorem implies that Ψ_p corresponds to $V_p^\sharp$ of this form, and thus has $R_p(\vec{N})$ eigenvalue $\varepsilon_p = -p^{-it_p}$, $T_p(p)$ eigenvalue $\varepsilon_p^2 = \chi(p)$, both $T_p(p, 1)$ and $T_p(1, p)$

eigenvalues $-\varepsilon_p$. (Note $T_p(1, p) = R_p(\vec{N})^{-1} T_p(p, 1) R_p(\vec{N})$.) Thus by the calculations of §1.3.2, Ψ_p has $T_p^{[p^n]}$ eigenvalues $\lambda_p^{[p^n]} = p^{-nit_p}$.

For $p \mid N^\chi$, π_p is equivalent to the pre-unitary continuous series $\pi(\tilde{\chi}_p | \cdot|_p^{it_p}, | \cdot|_p^{-it_p})$, with $t_p \in \mathbb{R}/\frac{2\pi}{\log(p)}\mathbb{Z}$, and the new vector V_p^χ has the form, for $\kappa \in \mathcal{O}_p^\times$,

$$\begin{aligned} V_p^\chi \left(\begin{bmatrix} \dot{a} & * \\ & \ddot{a} \end{bmatrix} \kappa \right) &= \tilde{\chi}_p(\dot{a}\kappa) \left| \frac{\dot{a}}{\ddot{a}} \right|_p^{\frac{1}{2}+it_p} V_p^\chi(1), \\ V_p^\chi \left(\begin{bmatrix} * & \dot{a} \\ & \ddot{a} \end{bmatrix} \kappa \right) &= 0. \end{aligned}$$

Therefore Ψ_p has $T_p^\chi(p^n, 1)$ eigenvalues $p^{\frac{n}{2}-nit_p}$, all $T_p^\chi(p^n)$ eigenvalues 1, and hence $T_p^\chi(1, p^n)$ eigenvalues $p^{\frac{n}{2}+nit_p}$.

For $p \mid d_B$, π_p is an irreducible representation of $D_p^\times$, trivial on $R_p^\times$, with unramified central character $\tilde{\chi}_p$. It is easy to see from this that π_p is one-dimensional and given by the character $|\nu(\cdot)|_p^{it_p}$, with $t_p \in \frac{i}{2} \log_p(\chi(p)) + \frac{\pi}{\log(p)}\mathbb{Z}/\frac{2\pi}{\log(p)}\mathbb{Z}$. Ψ_p must be a multiple of this character, and so has $T_p^{[p^n]}$ eigenvalues $\lambda_p^{[p^n]} = p^{-nit_p}$. The representation π_p^{JL} of $\text{GL}_2(\mathbb{Q}_p)$ associated to π_p by Jacquet–Langlands is the special representation $\sigma(| \cdot|_p^{\frac{1}{2}+it_p}, | \cdot|_p^{-\frac{1}{2}+it_p})$.

For $v = \infty$ and $k = 0$, we have $\tilde{\chi}_\infty = 1$, and π_∞ is equivalent to the pre-unitary continuous series $\pi(\text{sgn}^{\delta_\infty} | \cdot|_\infty^{s_\infty}, \text{sgn}^{\delta_\infty} | \cdot|_\infty^{-s_\infty})$. V_∞^0 corresponding to Ψ_∞ has the form,

$$V_\infty^0 \left(\begin{bmatrix} \dot{a} & * \\ & \ddot{a} \end{bmatrix} \kappa \right) = \text{sgn}^{\delta_\infty} \left(\frac{\dot{a}}{\ddot{a}} \right) \left| \frac{\dot{a}}{\ddot{a}} \right|_\infty^{\frac{1}{2}+s_\infty} V_\infty^0(1) \quad \text{for } \kappa \in \text{SO}(2, \mathbb{R}).$$

Thus Ψ_∞ has T_∞^- eigenvalue $(-1)^{\delta_\infty}$, T_∞^Δ eigenvalue $\lambda_\infty = (\frac{1}{4} - s_\infty^2) > 0$, and $T_{X_\infty}^{0,\xi} = T_{Y_\infty}^{0,\xi}$ eigenvalues

$$\begin{aligned} \lambda_\infty^{0,\xi} &= \int_{\mathbb{R}^+ \setminus B_\infty^+} (V_\infty^0(\beta)/V_\infty^0(1)) \phi_{X_\infty}^{0,\xi}(\beta) d_\infty^\times \beta \\ &= \int_{\mathbb{R}} \int_{\mathbb{R}^+} \int_0^{2\pi} y^{\frac{1}{2}+s_\infty} \frac{1}{\pi} \exp(-\pi\xi(y^2 + x^2 + 1)y^{-1}) y^{-2} dx dy d\theta \\ &= 2\xi^{-\frac{1}{2}} \int_{\mathbb{R}^+} \exp(-\pi\xi(y + y^{-1})) y^{s_\infty-1} dy \\ &= 4\xi^{-\frac{1}{2}} K_{s_\infty}(2\pi\xi). \end{aligned}$$

We are not considering $|k| = 1$, but we mention that in this case π_∞ is equivalent to the continuous series $\pi(\text{sgn}^{1+\delta_\infty} | \cdot|_\infty^{s_\infty}, \text{sgn}^{\delta_\infty} | \cdot|_\infty^{-s_\infty})$.

If $|k| \geq 2$, π_∞ is equivalent to the discrete series $\sigma(\text{sgn}^k | \cdot|_\infty^{s_\infty}, | \cdot|_\infty^{-s_\infty})$ for $s_\infty = \frac{|k|-1}{2}$. Upon restriction to $\text{SL}_2(\mathbb{R})$, this representation decomposes as the direct sum of the weight- $|k|$ holomorphic and anti-holomorphic discrete series, and the vector V_∞^k corresponding to Ψ_∞ is a lowest-weight vector in either the first or second of

these, depending on whether $k > 0$ or $k < 0$. Thus V_∞^k has the form,

$$V_\infty^k \left(\begin{bmatrix} \dot{a} & * \\ & \ddot{a} \end{bmatrix} \kappa \right) = \tilde{\chi}_\infty(\dot{a}\kappa) \left| \frac{\dot{a}}{\ddot{a}} \right|_\infty^{\frac{|k|}{2}} V_\infty^k(1) \quad \text{for } \kappa \in \text{SO}(2, \mathbb{R}),$$

and Ψ_∞ has T_∞^Δ eigenvalue $\lambda_\infty = \frac{|k|(2-|k|)}{4}$, while $T_\infty^- V_\infty^{\pm k} = V_\infty^{\mp k}$. We define

$$T_{X_\infty}^{k,\xi} \Psi_\infty = \lambda_{X_\infty}^{k,\xi} \Psi_\infty, \quad T_{Y_\infty}^{k,\xi} \Psi_\infty = \lambda_{Y_\infty}^{k,\xi} \Psi_\infty.$$

Our calculation is similar to that for $k = 0$, but we now use the mean value and conformal mapping properties of harmonic functions. These eigenvalues can also be computed by application of Selberg's lemma on invariant integral operators.

$$\begin{aligned} \lambda_{X_\infty}^{k,\xi} &= \int_{\mathbb{R}} \int_{\mathbb{R}^+} \int_0^{2\pi} y^{\frac{|k|}{2}} \frac{1}{\pi} \left(\frac{y+1-ix}{2y^{1/2}} \right)^k \exp \left(-\pi \xi \left(\frac{y^2+x^2+1}{y} \right) \right) y^{-2} dx dy d\theta \\ &= \int_{\mathbb{R}} \int_{\mathbb{R}^+} 2 \left(\frac{1-iz}{2} \right)^k \exp(-2\pi \xi \cosh \text{dist}(i, z)) y^{-2} dx dy \\ &= \int_{\mathbb{R}} \int_{\mathbb{R}^+} 2 \exp \left(-\pi \xi \left(\frac{y^2+x^2+1}{y} \right) \right) y^{-2} dx dy \\ &= 4 \xi^{-\frac{1}{2}} K_{-\frac{1}{2}}(2\pi \xi) = 2 \xi^{-1} e(i\xi), \\ \lambda_{Y_\infty}^{k,\xi} &= \int_{\mathbb{R}} \int_{\mathbb{R}^+} \int_0^{2\pi} y^{\frac{|k|}{2}} \frac{1}{\pi} \left(\frac{y-1-ix}{2y^{1/2}} \right)^k \exp \left(-\pi \xi \left(\frac{y^2+x^2+1}{y} \right) \right) y^{-2} dx dy d\theta \\ &= \int_{\mathbb{R}} \int_{\mathbb{R}^+} 2 \left(\frac{-1-iz}{2} \right)^k \exp(-2\pi \xi \cosh \text{dist}(i, z)) y^{-2} dx dy = 0. \end{aligned}$$

1.5 Whittaker Models

In this section, we restrict our attention to $B^\times = \text{GL}_2 = G$ (i.e. $d_B = 1$) and $\mathcal{O}(1, (1, M))$, M square-free. Let $e_\infty(x) = e(x) = \exp(2\pi i x)$ for $x \in \mathbb{R}$, and define the character e_p of $\mathbb{Q}_p$, with kernel $\mathbb{Z}_p$, on $x \in \mathbb{Z}[p^{-1}] \subset \mathbb{Q}_p$ by $e_p(x) = e(-x)$. Thus $e_{\mathbb{A}} = \prod_v e_v$ is trivial on $\mathbb{Q}$ and unramified.

Any $F \in L_0^2(G_{\mathbb{Q}} \backslash G_{\mathbb{A}}, \tilde{\chi})$ has a Fourier–Whittaker series expansion,

$$\begin{aligned} F(g) &= \sum_{\xi \in \mathbb{Q}^\times} \widehat{F} \left(\begin{bmatrix} \xi & \\ & 1 \end{bmatrix} g \right), \\ \widehat{F}(g) &:= \int_{\mathbb{Q} \backslash \mathbb{A}} F \left(\begin{bmatrix} 1 & x \\ & 1 \end{bmatrix} g \right) \overline{e_{\mathbb{A}}(x)} dx. \end{aligned}$$

We will see in §3.2.1 that

$$\langle F, F \rangle = \int_{\mathbb{A}^\times} \left| \widehat{F} \left(\begin{bmatrix} a & \\ & 1 \end{bmatrix} \right) \right|^2 |a|_{\mathbb{A}}^{-1} da.$$

Define the global Whittaker model $\mathcal{W} = \mathcal{W}(\pi) \simeq \pi$ as the image of π under $F \mapsto \widehat{F}$. Then $\mathcal{W} = \tilde{\otimes}_v \mathcal{W}_v$, where all $W_v \in \mathcal{W}_v \simeq \pi_v$ are smooth functions on $G_{\mathbb{Q}_v}$ s.t.

$$W_v \left(\begin{bmatrix} 1 & x \\ & 1 \end{bmatrix} g \right) = e_v(x) W_v(g) \quad \text{for } x \in \mathbb{Q}_v.$$

We will explicitly compute those W_v having the same right K_v types as forms in $\tilde{S}_k(1, (1, M), \tilde{\chi}, M^\flat)$. By the uniqueness of $\mathcal{W}_v$ proved in [17], it suffices for us to write down candidates W_v^* and check that they are $\mathcal{H}_v^*$ eigenvectors. Of course we could solve for these functions directly, by working backwards through the calculations below, but we find it easier to ‘guess’ them based on the results of §2.2.

An important feature of the Whittaker models is that they endow forms on GL_2 with an arithmetic normalization distinct from the spectral one. An eigenform F is called Hecke normalized if $\widehat{F} = \prod_v W_v^*$. The W_v^* have been scaled so that $W_p^*(1) = 1$ and W_∞^* has prescribed asymptotics in the cusp. We denote by $\check{W}_v^* \in \mathcal{W}_v^\vee$ the Whittaker functions of $\overline{F}$,

$$\check{W}_v^*(g) = \overline{W_v^*(\epsilon g)} \quad \text{for } \epsilon = \begin{bmatrix} -1 & \\ & 1 \end{bmatrix}.$$

For $p \nmid M$, we define W_p^0 by

$$W_p^0 \left(\begin{bmatrix} \dot{a} & \\ & \ddot{a} \end{bmatrix} \right) = \left(\frac{|\dot{a}p|^{\dot{s}} |\ddot{a}|^{\ddot{s}} - |\dot{a}p|^{\ddot{s}} |\ddot{a}|^{\dot{s}}}{|p|^{\dot{s}} - |p|^{\ddot{s}}} \right) |\frac{\dot{a}}{\ddot{a}}|^{\frac{1}{2}} \mathbb{I}_{\mathbb{Z}_p} \left(\frac{\dot{a}}{\ddot{a}} \right),$$

and check that

$$\begin{aligned} T_p(p, 1) W_p^0 \left(\begin{bmatrix} a & \\ & 1 \end{bmatrix} \right) &= \sum_{x \bmod p} W_p^0 \left(\begin{bmatrix} ap & ax \\ & 1 \end{bmatrix} \right) + W_p^0 \left(\begin{bmatrix} a & \\ & p \end{bmatrix} \right) \\ &= \sum_{x \bmod p} e_p(ax) \left(\frac{|ap^2|^{\dot{s}} - |ap^2|^{\ddot{s}}}{|p|^{\dot{s}} - |p|^{\ddot{s}}} \right) |ap|^{\frac{1}{2}} \mathbb{I}_{\mathbb{Z}_p}(ap) \\ &\quad + \left(\frac{|ap|^{\dot{s}} |p|^{\ddot{s}} - |ap|^{\ddot{s}} |p|^{\dot{s}}}{|p|^{\dot{s}} - |p|^{\ddot{s}}} \right) |\frac{a}{p}|^{\frac{1}{2}} \mathbb{I}_{\mathbb{Z}_p} \left(\frac{a}{p} \right) \\ &= \left(\frac{p^{-\dot{s}} |ap|^{\dot{s}} - p^{-\ddot{s}} |ap|^{\ddot{s}}}{p^{-\dot{s}} - p^{-\ddot{s}}} \right) p^{\frac{1}{2}} |a|^{\frac{1}{2}} \mathbb{I}_{\mathbb{Z}_p}(a) \\ &\quad + \left(\frac{p^{-\ddot{s}} |ap|^{\dot{s}} - p^{-\dot{s}} |ap|^{\ddot{s}}}{p^{-\dot{s}} - p^{-\ddot{s}}} \right) p^{\frac{1}{2}} |a|^{\frac{1}{2}} \mathbb{I}_{\mathbb{Z}_p}(a) \\ &= \lambda_p W_p^0 \left(\begin{bmatrix} a & \\ & 1 \end{bmatrix} \right). \end{aligned}$$

Recall that the case of $p \mid M^\flat$ is essentially the same as the previous one, but we must consider a two-dimensional space of oldvectors. Taking W_p^0 as above, we have

already shown $\dot{W}_p^0 \in \mathcal{W}_p$, and we simply define

$$\begin{aligned}\ddot{W}_p^0 &= \begin{bmatrix} 1 & \\ & p \end{bmatrix} \dot{W}_p^0 = R_p(1, p) \dot{W}_p^0, \\ W_p^b &= \dot{W}_p^0 + \overline{\varepsilon}_p \ddot{W}_p^0, \quad \varepsilon_p^2 = \chi(p).\end{aligned}$$

For $p \mid M^\sharp$, we define

$$\begin{aligned}W_p^\sharp \left(\begin{bmatrix} \dot{a} & \\ & \ddot{a} \end{bmatrix} \right) &= |\dot{a}\ddot{a}|_p^{\text{it}_p} \left| \frac{\dot{a}}{\ddot{a}} \right|_p \mathbb{I}_{\mathbb{Z}_p} \left(\frac{\dot{a}}{\ddot{a}} \right), \\ W_p^\sharp \left(\begin{bmatrix} & \dot{a} \\ \ddot{a} & \end{bmatrix} \right) &= -|\dot{a}\ddot{a}|_p^{\text{it}_p} \left| \frac{\dot{a}p}{\ddot{a}} \right|_p \mathbb{I}_{\mathbb{Z}_p} \left(\frac{\dot{a}p}{\ddot{a}} \right),\end{aligned}$$

and check that

$$\begin{aligned}R_p(1, p)[W_p^\sharp] \left(\begin{bmatrix} a & \\ & 1 \end{bmatrix} \right) &= W_p^\sharp \left(\begin{bmatrix} & a \\ p & \end{bmatrix} \right) \\ &= -p^{-\text{it}_p} W_p^\sharp \left(\begin{bmatrix} a & \\ & 1 \end{bmatrix} \right), \\ R_p(1, p)[W_p^\sharp] \left(\begin{bmatrix} & a \\ p & \end{bmatrix} \right) &= p^{-2\text{it}_p} W_p^\sharp \left(\begin{bmatrix} a & \\ & 1 \end{bmatrix} \right) \\ &= -p^{-\text{it}_p} W_p^\sharp \left(\begin{bmatrix} & a \\ p & \end{bmatrix} \right), \\ T_p(p, 1)[W_p^\sharp] \left(\begin{bmatrix} a & \\ & 1 \end{bmatrix} \right) &= \sum_{x \bmod p} W_p^\sharp \left(\begin{bmatrix} ap & ax \\ & 1 \end{bmatrix} \right) \\ &= \sum_{x \bmod p} \mathbf{e}_p(ax) |ap|_p^{1+\text{it}_p} \mathbb{I}_{\mathbb{Z}_p}(ap) \\ &= |ap|_p^{1+\text{it}_p} p \mathbb{I}_{\mathbb{Z}_p}(a) = p^{-\text{it}_p} W_p^\sharp \left(\begin{bmatrix} a & \\ & 1 \end{bmatrix} \right), \\ T_p(p, 1)[W_p^\sharp] \left(\begin{bmatrix} & a \\ 1 & \end{bmatrix} \right) &= \sum_{x \bmod p} W_p^\sharp \left(\begin{bmatrix} & a \\ p & x \end{bmatrix} \right) \\ &= W_p^\sharp \left(\begin{bmatrix} & a \\ p & \end{bmatrix} \right) + \sum_{x \not\equiv 0} W_p^\sharp \left(\begin{bmatrix} 1 & \frac{a}{x} \\ & 1 \end{bmatrix} \begin{bmatrix} ap & \\ & 1 \end{bmatrix} \begin{bmatrix} -\frac{1}{x} & \\ p & x \end{bmatrix} \right) \\ &= -p |ap|_p^{1+\text{it}_p} \mathbb{I}_{\mathbb{Z}_p}(a) + \sum_{x \not\equiv 0} \mathbf{e}_p\left(\frac{a}{x}\right) |ap|_p^{1+\text{it}_p} \mathbb{I}_{\mathbb{Z}_p}(ap) \\ &= -p |ap|_p^{1+\text{it}_p} \mathbb{I}_{\mathbb{Z}_p}(a) + |ap|_p^{1+\text{it}_p} (p \mathbb{I}_{\mathbb{Z}_p}(a) - \mathbb{I}_{\mathbb{Z}_p}(ap)) \\ &= -|ap|_p^{1+\text{it}_p} \mathbb{I}_{\mathbb{Z}_p}(ap) = p^{-\text{it}_p} W_p^\sharp \left(\begin{bmatrix} & a \\ 1 & \end{bmatrix} \right).\end{aligned}$$

For $p \mid M^\chi$, we define

$$\begin{aligned} W_p^\chi \left(\begin{bmatrix} \dot{a} & \\ & \ddot{a} \end{bmatrix} \right) &= \tilde{\chi}_p(\dot{a}) \left| \frac{\dot{a}}{\ddot{a}} \right|_p^{\frac{1}{2}+it_p} \mathbb{I}_{\mathbb{Z}_p} \left(\frac{\dot{a}}{\ddot{a}} \right), \\ W_p^\chi \left(\begin{bmatrix} & \dot{a} \\ \ddot{a} & \end{bmatrix} \right) &= \overline{\mathcal{G}_p} \tilde{\chi}_p(\ddot{a}) \left| \frac{\dot{a}p^2}{\ddot{a}} \right|_p^{\frac{1}{2}-it_p} \mathbb{I}_{\mathbb{Z}_p} \left(\frac{\dot{a}p}{\ddot{a}} \right), \\ \text{for } \mathcal{G}_p &= \int_{\frac{1}{p}\mathbb{Z}_p^\times} \overline{\tilde{\chi}_p(x)} e_p(x) dx, \end{aligned}$$

and check that

$$\begin{aligned} T_p^\chi(p, 1)[W_p^\chi] \left(\begin{bmatrix} a & \\ & 1 \end{bmatrix} \right) &= \sum_{x \bmod p} W_p^\chi \left(\begin{bmatrix} ap & ax \\ & 1 \end{bmatrix} \right) \overline{\tilde{\chi}_p(p)} \\ &= \sum_{x \bmod p} e_p(ax) \tilde{\chi}_p(a) |ap|_p^{\frac{1}{2}+it_p} \mathbb{I}_{\mathbb{Z}_p}(ap) \\ &= \tilde{\chi}_p(a) |ap|_p^{\frac{1}{2}+it_p} p \mathbb{I}_{\mathbb{Z}_p}(a) = p^{\frac{1}{2}-it_p} W_p^\chi \left(\begin{bmatrix} a & \\ & 1 \end{bmatrix} \right), \\ T_p^\chi(p, 1)[W_p^\chi] \left(\begin{bmatrix} & a \\ 1 & \end{bmatrix} \right) &= \sum_{x \bmod p} W_p^\chi \left(\begin{bmatrix} & a \\ p & x \end{bmatrix} \right) \overline{\tilde{\chi}_p(p)} \\ &= W_p^\chi \left(\begin{bmatrix} & a \\ p & \end{bmatrix} \right) \frac{1}{p} + \sum_{x \neq 0} W_p^\chi \left(\begin{bmatrix} 1 & \frac{a}{x} \\ & 1 \end{bmatrix} \begin{bmatrix} ap & \\ & 1 \end{bmatrix} \begin{bmatrix} -\frac{1}{x} & \\ p & x \end{bmatrix} \right) \frac{1}{p} \\ &= \overline{\mathcal{G}_p} |ap|_p^{\frac{1}{2}-it_p} \mathbb{I}_{\mathbb{Z}_p}(a) + \sum_{x \neq 0} e_p\left(\frac{a}{x}\right) \tilde{\chi}_p\left(-\frac{a}{x}\right) |ap|_p^{\frac{1}{2}+it_p} \mathbb{I}_{\mathbb{Z}_p}(ap) \\ &= \overline{\mathcal{G}_p} |ap|_p^{\frac{1}{2}-it_p} \mathbb{I}_{\mathbb{Z}_p}(a) + \overline{\mathcal{G}_p} \mathbb{I}_{\mathbb{Z}_p^\times}(ap) = p^{\frac{1}{2}-it_p} W_p^\chi \left(\begin{bmatrix} & a \\ 1 & \end{bmatrix} \right). \end{aligned}$$

For $v = \infty$, recall the classical Whittaker function $w_{\kappa, \mu}$ which uniquely solves

$$\begin{aligned} w_{\kappa, \mu}''(y) + \left(-\frac{1}{4} + \frac{\kappa}{y} + \frac{\frac{1}{4} - \mu^2}{y^2} \right) w_{\kappa, \mu}(y) &= 0 \\ \text{under } w_{\kappa, \mu}(y) &\sim y^\kappa e^{-\frac{y}{2}} \text{ as } y \rightarrow \infty. \end{aligned}$$

In particular,

$$\begin{aligned} w_{\kappa, \mu}(y) &= \frac{y^{\mu+\frac{1}{2}} e^{-\frac{y}{2}}}{\Gamma(\mu - \kappa + \frac{1}{2})} \int_0^\infty e^{-yt} t^{\mu-\kappa-\frac{1}{2}} (1+t)^{\mu+\kappa-\frac{1}{2}} dt, \\ w_{0, \mu}(y) &= \left(\frac{y}{\pi} \right)^{\frac{1}{2}} K_\mu\left(\frac{y}{2}\right), \quad w_{\frac{k}{2}, \frac{k-1}{2}}(y) = y^{\frac{k}{2}} e^{-\frac{y}{2}}. \end{aligned}$$

If $k = 0$, define $W_\infty^0 \in \mathcal{W}_\infty$ by

$$W_\infty^0 \left(\begin{bmatrix} \dot{a} & \\ & \ddot{a} \end{bmatrix} \right) = \operatorname{sgn}^{\delta_\infty} \left(\frac{\dot{a}}{\ddot{a}} \right) w_{0,s_\infty} \left(\frac{\dot{a}}{\ddot{a}} \right).$$

This has moderate growth and satisfies

$$\begin{aligned} T_\infty^- [W_\infty^0] &= (-1)^{\delta_\infty} W_\infty^0, \\ T_\infty^\Delta [W_\infty^0] &= \left(\frac{1}{4} - s_\infty^2 \right) W_\infty^0. \end{aligned}$$

If $|k| \geq 2$, define $W_\infty^k \in \mathcal{W}_\infty$ by

$$W_\infty^k \left(\begin{bmatrix} \dot{a} & \\ & \ddot{a} \end{bmatrix} \right) = \mathbb{I}_{k\mathbb{R}^+} \left(\frac{\dot{a}}{\ddot{a}} \right) w_{\frac{|k|}{2}, \frac{|k|-1}{2}} \left(\frac{\dot{a}}{\ddot{a}} \right),$$

which has moderate growth and satisfies, for $\pm k \geq 2$,

$$\begin{aligned} R_k^\mp [W_\infty^k] &= 0, \\ T_\infty^\Delta [W_\infty^k] &= \frac{|k|(2-|k|)}{4} W_\infty^k. \end{aligned}$$

Chapter 2

Theta Lifting

2.1 Basic Setup

2.1.1 Groups

Following [8], let $G = \mathrm{GL}_2$, $H = \mathrm{GSp}_6$, $\mathbf{G} = G^3 \cap H$ as linear algebraic groups over $\mathbb{Q}$. The similitude character of H and its restriction to $\mathbf{G}$ will be denoted ν (and distinguished by context from the reduced norm of B). Also consider B , $B^\times$, $H' = \mathrm{GO}(B, \nu)$, $\mathbf{G}' = H'^3 \cap \mathrm{GO}(B^3, \oplus^3 \nu)$ as linear algebraic groups over $\mathbb{Q}$, again denoting the similitude characters by ν . Write $\check{H}'$ for the connected component of the identity in H' , and define $\rho : B^\times \times B^\times \rightarrow \check{H}'$ by $\rho(\dot{\alpha}, \ddot{\alpha})(\beta) = \dot{\alpha}\beta\ddot{\alpha}^{-1}$. We see there is an exact sequence

$$1 \longrightarrow Z(B^\times) \xrightarrow{\mathrm{diag}} (B^\times \times B^\times) \xrightarrow{\rho} \check{H}' \longrightarrow 1.$$

We may view the anti-involution ι as an element of order two in $H'_\mathbb{Q}$, $\iota(\beta) = \beta^\iota$. It and $\check{H}'$ generate $H' = \check{H}' \rtimes \langle \iota \rangle$ as a semi-direct product, $\rho(\dot{\alpha}, \ddot{\alpha})\iota = \iota\rho\left(\frac{\ddot{\alpha}}{\nu(\ddot{\alpha})}, \frac{\dot{\alpha}}{\nu(\dot{\alpha})}\right)$. Note that

$$\begin{aligned} \prod_v (\mathbb{Z}/2\mathbb{Z}) &\xrightarrow{\sim} \langle \iota \rangle_\mathbb{A}, \\ (\delta_v) &\mapsto (\iota_v^{\delta_v}). \end{aligned}$$

To compute inner-products on H' , we will use the parametrization

$$\rho : PB^\times \times PB^\times \xrightarrow{\sim} P\check{H}'.$$

We will also need Shimizu's parametrization,

$$\begin{aligned} B^\times \rtimes PB^\times &\xrightarrow{\sim} \check{H}', \\ B^{(\delta)} \times PB^\times &\xrightarrow{1-1} \check{H}'^{(\delta)}, \\ (\dot{\beta}, \ddot{\beta}) &\longmapsto \rho(\dot{\beta}\ddot{\beta}, \ddot{\beta}). \end{aligned}$$

2.1.2 Measures

We now describe the canonical Tamagawa measures on each of these groups, as in [42, 40]. Let $dx = dx_v$ be the Haar measure on $\mathbb{Q}_v$ such that the Fourier transform is self-dual ($\text{vol}(\mathbb{R}/\mathbb{Z}) = \text{vol}(\mathbb{Z}_p) = 1 = \text{vol}(\mathbb{A}/\mathbb{Q})$): For $\varphi \in \mathcal{S}(\mathbb{Q}_v)$,

$$\mathcal{F}\varphi(y) = \int_{\mathbb{Q}_v} \varphi(y) e_v(xy) dx = \mathcal{F}^{-1}\varphi(-y).$$

Then define $d^\times x_v = \frac{\zeta_v(1)}{|x|_v} dx_v$, so that $\text{vol}^\times(\mathbb{R}^+/e^\mathbb{Z}) = \text{vol}^\times(\mathbb{Z}_p^\times) = 1$.

Identify B_v with its algebraic dual B_v^* using the non-degenerate symmetric bilinear form $\langle \alpha, \beta \rangle = \text{tr}(\alpha\beta^t) = \alpha\beta^t + \beta\alpha^t$. Now choose the Haar measure $d\alpha = d\alpha_v$ on B_v so that the ‘ ι -twisted’ Fourier transform is self-dual: For $\varphi \in \mathcal{S}(B_v)$,

$$\mathcal{F}\varphi(\beta) = \int_{B_v} \varphi(\alpha) e_v(\langle \alpha, \beta \rangle) d\alpha = \mathcal{F}^{-1}\varphi(-\beta).$$

If $v \nmid d_B$, $d\alpha = \prod_{ij} d\alpha_{ij}$, while for $p \mid d_B$, $\text{vol}(R_p) = \frac{1}{p}$. It follows from calculating $\text{vol}(B_\infty/\mathcal{O}(d_B, \vec{N})) = d_B N$ that $\text{vol}(B_\mathbb{A}/B_\mathbb{Q}) = 1$.

As before, define $d^\times \alpha_v = \frac{\zeta_v(1)}{|\alpha|_v} d\alpha_v$. It is well known and follows from the calculations in §1.3 that

$$\text{vol}^\times(\mathcal{O}_p^\times(d_B, \vec{N})) = \zeta_p(2)^{-1} \begin{cases} (p-1)^{-1} & \text{if } p \mid d_B, \\ (p+1)^{-1} & \text{if } p \mid N, \\ 1 & \text{if } p \nmid d_B N. \end{cases}$$

On B_∞ , $d^\times \alpha_\infty = d^\times a_1 d^\times a_2 dx d\theta$ in terms of the coordinates

$$\alpha_\infty = \begin{bmatrix} a_1 & \\ & a_2 \end{bmatrix} \begin{bmatrix} 1 & x \\ & 1 \end{bmatrix} \kappa_\theta, \quad \text{for } 0 \leq \theta < \pi.$$

We will also write $d^\times \alpha_v$ for the measure on $PB_v^\times$ compatible with the exact sequence

$$1 \longrightarrow \mathbb{Q}_v^\times \longrightarrow B_v^\times \longrightarrow PB_v^\times \longrightarrow 1,$$

and $d^{(1)}\alpha_v$ for the measure on $B_v^{(1)}$ compatible with

$$1 \longrightarrow B_v^{(1)} \longrightarrow B_v^\times \xrightarrow{\nu} \mathbb{Q}_v^\times \longrightarrow 1.$$

To be explicit,

$$\text{vol}^\times(P\mathcal{O}_p^\times(d_B, \vec{N})) = \text{vol}^\times(\mathcal{O}_p^\times(d_B, \vec{N})),$$

while $d^\times \alpha_\infty = d^\times a \, dx \, d\theta$ in the $PB_\infty^\times$ coordinates

$$\alpha_\infty = \begin{bmatrix} a & \\ & 1 \end{bmatrix} \begin{bmatrix} 1 & x \\ & 1 \end{bmatrix} \kappa_\theta, \quad \text{for } 0 \leq \theta < \pi.$$

Also

$$\text{vol}^{(1)}(\mathcal{O}_p^{(1)}(d_B, \vec{N})) = \text{vol}^\times(\mathcal{O}_p^\times(d_B, \vec{N})),$$

while $d^{(1)}\alpha_\infty = d^\times a \, dx \, d\theta$ in the $B_\infty^{(1)}$ coordinates

$$\alpha_\infty = \begin{bmatrix} a & \\ & a^{-1} \end{bmatrix} \begin{bmatrix} 1 & x \\ & 1 \end{bmatrix} \kappa_\theta, \quad \text{for } 0 \leq \theta < \pi.$$

It is well known (see [40]) that $\text{vol}^\times(PB_\mathbb{Q}^\times \backslash PB_\mathbb{A}^\times) = 2$ and $\text{vol}^{(1)}(B_\mathbb{Q}^{(1)} \backslash B_\mathbb{A}^{(1)}) = 1$. The difference between $d^\times \alpha_\infty$ and $d^{(1)}\alpha_\infty$ may be seen as follows: Lift $d^\times \alpha_\infty$ to the double cover $\mathbb{R}^+ \backslash B_\infty^\times$ of $PB_\infty^\times$. Then $\frac{1}{2} d^\times \alpha_\infty$ is compatible with

$$1 \longrightarrow B_\infty^{(1)} \longrightarrow \mathbb{R}^+ \backslash B_\infty^\times \xrightarrow{\nu} \mathbb{R}^+ \backslash \mathbb{R}^\times \longrightarrow 1.$$

The Tamagawa measures on $G = \text{GL}_2$, $PG = \text{PGL}_2$, $G^{(1)} = \text{SL}_2$ are special cases of those above, for $d_B = 1$. Now define $d\mathbf{g}_v$ on $\mathbf{G}_v$ to be compatible with

$$1 \longrightarrow (G_v^{(1)})^3 \longrightarrow \mathbf{G}_v \xrightarrow{\nu} \mathbb{Q}_v^\times \longrightarrow 1.$$

Explicitly,

$$\text{vol}(\mathcal{O}_p^\times(1, \vec{N})^3 \cap \mathbf{G}_p) = \text{vol}^\times(\mathcal{O}_p^\times(1, \vec{N}))^3,$$

and $d\mathbf{g}_\infty = d^\times a \prod_j (d^\times a_j \, dx_j \, d\theta_j)$ in the $\mathbf{G}_\infty$ coordinates

$$\mathbf{g}_{\infty j} = \begin{bmatrix} a & \\ & 1 \end{bmatrix} \begin{bmatrix} a_j & \\ & a_j^{-1} \end{bmatrix} \begin{bmatrix} 1 & x_j \\ & 1 \end{bmatrix} \kappa_{\theta_j}, \quad \text{for } 0 \leq \theta_j < \pi.$$

Also define $d\mathbf{g}_v$ on $P\mathbf{G}_v$ to be compatible with

$$1 \longrightarrow \mathbb{Q}_v^\times \longrightarrow \mathbf{G}_v \longrightarrow P\mathbf{G}_v \longrightarrow 1.$$

Then

$$\text{vol}(P(\mathcal{O}_p^\times(1, \vec{N})^3 \cap \mathbf{G}_p)) = \text{vol}^\times(\mathcal{O}_p^\times(1, \vec{N}))^3,$$

and $d\mathbf{g}_\infty = 2 \prod_j (d^\times a_j dx_j d\theta_j)$ in the $P\mathbf{G}_\infty$ coordinates

$$\mathbf{g}_{\infty j} = \begin{bmatrix} \epsilon & \\ & 1 \end{bmatrix} \begin{bmatrix} a_j & \\ & a_j^{-1} \end{bmatrix} \begin{bmatrix} 1 & x_j \\ & 1 \end{bmatrix} \kappa_{\theta_j},$$

for $\epsilon^2 = 1$, $a_1 a_2 a_3 > 0$, $0 \leq \theta_j < \pi$.

It is easy to check that $\text{vol}(P\mathbf{G}_\mathbb{Q} \backslash P\mathbf{G}_\mathbb{A}) = 2$.

2.1.3 Dual-Pairs

Extend the definition of $\langle \cdot, \cdot \rangle$ to B_v^n as an orthogonal direct sum, and define $\mathcal{F}$, $d\alpha_v$ on B_v^n as before. We may write $\mathcal{F}_j$ for the Fourier transform in β_j alone, $\beta = (\beta_j) \in B_v^n$. The Weil representation ω of $\text{Sp}_{2n}(\mathbb{Q}_v)$ on $\mathcal{S}(B_v^n)$ is uniquely determined by: [17, 34]

$$\begin{aligned} \omega \left(\begin{bmatrix} 1_n & U \\ & 1_n \end{bmatrix} \right) \varphi(\beta) &= e_v(\tfrac{1}{2} \langle U\beta, \beta \rangle) \varphi(\beta) \quad \text{for } U = U^t \in M_n(\mathbb{Q}_v), \\ \omega \left(\begin{bmatrix} A^t & \\ & A^{-1} \end{bmatrix} \right) \varphi(\beta) &= |\det A|_v^2 \varphi(A\beta) \quad \text{for } A \in \text{GL}_n(\mathbb{Q}_v), \\ \omega \left(\begin{bmatrix} & 1_n \\ -1_n & \end{bmatrix} \right) \varphi(\beta) &= (-1)^{n\mathbb{I}_v|d_B} \mathcal{F}\varphi(\beta). \end{aligned}$$

Now consider the similitude dual-pairs

$$\begin{aligned} R(X, Y) &= \{(x, y) \in X \times Y; \nu(x) = \nu(y)\}, \\ \text{for } (n, X, Y) &= (1, G, H'), (3, \mathbf{G}, \mathbf{G}'), (3, H, H'). \end{aligned}$$

We extend the definition of ω to each of these as follows: Let L denote the unitary left regular representation of Y_v on $\mathcal{S}(B_v^n)$,

$$\begin{aligned} L(h') \varphi(\beta) &= |\nu(h')|_v^{-1} \varphi((h')^{-1}\beta), \\ L(\mathbf{g}') \varphi(\beta) &= |\nu(\mathbf{g}')|_v^{-3} \varphi((h'_j)^{-1}\beta_j), \\ L(h') \varphi(\beta) &= |\nu(h')|_v^{-3} \varphi((h')^{-1}\beta_j). \end{aligned}$$

If $\nu(x, y) = \delta$, set $\alpha = \begin{bmatrix} 1_n & \\ & \delta 1_n \end{bmatrix} \in X_v$, $x^{(1)} = x\alpha^{-1}$, ${}^{(1)}x = \alpha^{-1}x$. Then

$$\omega(x, y) = \omega(x^{(1)}) L(y) = L(y) \omega({}^{(1)}x)$$

defines a representation of $R(X, Y)_v$ on $\mathcal{S}(B_v^n)$.

Note $\omega(z, z) = 1$ for $z \in \mathbb{Q}_v^\times$, since $\omega \left(\begin{bmatrix} z 1_n & \\ & z^{-1} 1_n \end{bmatrix} \right) = L(z)^{-1}$.

2.2 Jacquet–Langlands Correspondence

In this section we explicitly compute the Shimizu theta lift and its adjoint, which realize the Jacquet–Langlands correspondence.

2.2.1 Shimizu’s Theta Lift

For $\dot{\Psi}, \ddot{\Psi}$ as in §1.2 and $\eta \in \prod_v (\mathbb{Z}/2\mathbb{Z})$, define $\check{F}' \in L_0^2(\check{H}'_{\mathbb{Q}} \backslash \check{H}'_{\mathbb{A}}, \check{\chi})$ by

$$\check{F}'(\rho(\dot{\beta}, \ddot{\beta})) = \dot{\Psi}^{\eta}(\dot{\beta}) \overline{\ddot{\Psi}^{\eta}(\ddot{\beta})},$$

with matching local data $\check{\varphi} \in \mathcal{S}(B_{\mathbb{A}})$ as in §2.3. Shimizu’s theta lift is defined in [34]: For $g \in G_{\mathbb{A}}^{(\delta)}$,

$$\begin{aligned} \check{\Theta}'_{\check{\varphi}}(\check{F}')(g) &= \frac{1}{2} \int_{\check{H}'_{\mathbb{Q}}^{(1)} \backslash \check{H}'_{\mathbb{A}}^{(\delta)}} \sum_{\alpha \in B_{\mathbb{Q}}} \omega(g, h') \check{\varphi}(\alpha) \check{F}'(h') dh' \\ &= \frac{1}{2} \int_{PB_{\mathbb{Q}}^{\times} \backslash PB_{\mathbb{A}}^{\times}} \int_{B_{\mathbb{Q}}^{(1)} \backslash B_{\mathbb{A}}^{(\delta)}} \sum_{\alpha \in B_{\mathbb{Q}}} \omega(g, \rho(\dot{\beta}\ddot{\beta}, \ddot{\beta})) \check{\varphi}(\alpha) \\ &\quad \cdot \dot{\Psi}^{\eta}(\dot{\beta}\ddot{\beta}) \overline{\ddot{\Psi}^{\eta}(\ddot{\beta})} d^{(1)}\dot{\beta} d^{\times}\ddot{\beta} \\ &= \frac{1}{2} \int_{PB_{\mathbb{Q}}^{\times} \backslash PB_{\mathbb{A}}^{\times}} \int_{B_{\mathbb{Q}}^{(1)} \backslash B_{\mathbb{A}}^{(\delta)}} \sum_{\alpha \in B_{\mathbb{Q}}} |\delta|_{\mathbb{A}}^{-1} \omega^{(1)}(g) \check{\varphi}((\dot{\beta}\ddot{\beta})^{-1} \alpha \ddot{\beta}) \\ &\quad \cdot \dot{\Psi}^{\eta}(\dot{\beta}\ddot{\beta}) \overline{\ddot{\Psi}^{\eta}(\ddot{\beta})} d^{(1)}\dot{\beta} d^{\times}\ddot{\beta}. \end{aligned}$$

This expression is readily converted into a Fourier series, following [34]. First, decompose the sum over $B_{\mathbb{Q}}$ into left $B_{\mathbb{Q}}^{(1)}$ orbits. Note that $B_{\mathbb{Q}}^{(1)} \backslash B_{\mathbb{Q}}^{\times} \simeq \mathbb{Q}^{\times}$ by ν , and if $B_{\mathbb{Q}}$ is a division algebra, $B_{\mathbb{Q}}^{(0)} = \{0\}$. In the unramified case $B = M$, the remaining elements are those $\alpha \in M_{\mathbb{Q}}$ of rank 1, and every such can be written as $\alpha = \gamma^{-1} \tilde{\alpha}$, for representatives of unique classes $\tilde{\alpha} \in \mathbb{P} \begin{bmatrix} \mathbb{Q} & \mathbb{Q} \\ 0 & 0 \end{bmatrix}$, $\gamma \in N_{\mathbb{Q}} \backslash \mathrm{SL}_2(\mathbb{Q})$.

The contribution to $\check{\Theta}'_{\check{\varphi}}(\check{F}')$ from $\alpha = 0$ is proportional to

$$\int_{PB_{\mathbb{Q}}^{\times} \backslash PB_{\mathbb{A}}^{\times}} \left(\int_{B_{\mathbb{Q}}^{(1)} \backslash B_{\mathbb{A}}^{(\delta)}} \dot{\Psi}^{\eta}(\dot{\beta}\ddot{\beta}) d^{(1)}\dot{\beta} \right) \overline{\ddot{\Psi}^{\eta}(\ddot{\beta})} d^{\times}\ddot{\beta} = 0,$$

since $\dot{\pi}$ is not one-dimensional, by the argument in [34]. For $B = M$ (which is not considered in [34], but see [36]), the contribution from all $\alpha \in M_{\mathbb{Q}}^{(0)} \setminus \{0\}$ equals

$$\begin{aligned} \frac{1}{2} \int_{PG_{\mathbb{Q}} \backslash PG_{\mathbb{A}}} \int_{G_{\mathbb{Q}}^{(1)} \backslash G_{\mathbb{A}}^{(\delta)}} \sum_{\tilde{\alpha}} \sum_{\gamma \in N_{\mathbb{Q}} \backslash G_{\mathbb{Q}}^{(1)}} \\ |\delta|_{\mathbb{A}}^{-1} \omega^{(1)}(g) \check{\varphi}((\gamma \dot{\beta} \ddot{\beta})^{-1} \tilde{\alpha} \ddot{\beta}) \dot{\Psi}^{\eta}(\dot{\beta}\ddot{\beta}) \overline{\ddot{\Psi}^{\eta}(\ddot{\beta})} d^{(1)}\dot{\beta} d^{\times}\ddot{\beta} \end{aligned}$$

$$\begin{aligned}
&= \sum_{\tilde{\alpha}} \frac{1}{2} \int_{PG_{\mathbb{Q}} \backslash PG_{\mathbb{A}}} \int_{N_{\mathbb{Q}} \backslash G_{\mathbb{A}}^{(\delta)}} \\
&\quad |\delta|_{\mathbb{A}}^{-1} \omega^{(1)}(g) \check{\varphi}((\dot{\beta}\ddot{\beta})^{-1} \tilde{\alpha}\ddot{\beta}) \dot{\Psi}^{\eta}(\dot{\beta}\ddot{\beta}) \overline{\ddot{\Psi}^{\eta}(\ddot{\beta})} d^{(1)}\dot{\beta} d^{\times}\ddot{\beta} \\
&= \sum_{\tilde{\alpha}} \frac{1}{2} \int_{PG_{\mathbb{Q}} \backslash PG_{\mathbb{A}}} \int_{N_{\mathbb{A}} \backslash G_{\mathbb{A}}^{(\delta)}} |\delta|_{\mathbb{A}}^{-1} \omega^{(1)}(g) \check{\varphi}((\dot{\beta}\ddot{\beta})^{-1} \tilde{\alpha}\ddot{\beta}) \\
&\quad \cdot \left(\int_{N_{\mathbb{Q}} \backslash N_{\mathbb{A}}} \dot{\Psi}^{\eta}(n\dot{\beta}\ddot{\beta}) dn \right) \overline{\ddot{\Psi}^{\eta}(\ddot{\beta})} d^{(1)}\dot{\beta} d^{\times}\ddot{\beta} = 0,
\end{aligned}$$

since $\dot{\pi}$ is cuspidal. Therefore in all cases under consideration,

$$\begin{aligned}
\check{\Theta}'_{\check{\varphi}}(\check{F}')(g) &= \frac{1}{2} \int_{PB_{\mathbb{Q}}^{\times} \backslash PB_{\mathbb{A}}^{\times}} \int_{B_{\mathbb{Q}}^{(1)} \backslash B_{\mathbb{A}}^{(\delta)}} \sum_{\tilde{\alpha} \in B_{\mathbb{Q}}^{(1)} \backslash B_{\mathbb{Q}}^{\times}} \sum_{\gamma \in B_{\mathbb{Q}}^{(1)}} \\
&\quad \omega(g, \rho(\gamma\dot{\beta}\ddot{\beta}, \ddot{\beta})) \check{\varphi}(\tilde{\alpha}) \dot{\Psi}^{\eta}(\dot{\beta}\ddot{\beta}) \overline{\ddot{\Psi}^{\eta}(\ddot{\beta})} d^{(1)}\dot{\beta} d^{\times}\ddot{\beta} \\
&= \sum_{\tilde{\alpha} \in B_{\mathbb{Q}}^{(1)} \backslash B_{\mathbb{Q}}^{\times}} \frac{1}{2} \int_{PB_{\mathbb{Q}}^{\times} \backslash PB_{\mathbb{A}}^{\times}} \int_{B_{\mathbb{A}}^{(\delta)}} \\
&\quad \omega(g, \rho(\dot{\beta}\ddot{\beta}, \ddot{\beta})) \check{\varphi}(\tilde{\alpha}) \dot{\Psi}^{\eta}(\dot{\beta}\ddot{\beta}) \overline{\ddot{\Psi}^{\eta}(\ddot{\beta})} d^{(1)}\dot{\beta} d^{\times}\ddot{\beta} \\
&= \sum_{\tilde{\alpha} \in B_{\mathbb{Q}}^{(1)} \backslash B_{\mathbb{Q}}^{\times}} \frac{1}{2} \int_{PB_{\mathbb{Q}}^{\times} \backslash PB_{\mathbb{A}}^{\times}} \int_{B_{\mathbb{A}}^{(\delta)}} \omega \left(\begin{bmatrix} \nu(\tilde{\alpha}) & \\ & 1 \end{bmatrix}, \rho(\tilde{\alpha}', 1) \right) \circ \\
&\quad \omega(g, \rho(\dot{\beta}\ddot{\beta}, \ddot{\beta})) \check{\varphi}(1) \dot{\Psi}^{\eta}(\dot{\beta}\ddot{\beta}) \overline{\ddot{\Psi}^{\eta}(\ddot{\beta})} d^{(1)}\dot{\beta} d^{\times}\ddot{\beta} \\
&= \sum_{\xi \in \mathbb{Q}^{\times}} \widehat{\check{\Theta}'_{\check{\varphi}}(\check{F}')} \left(\begin{bmatrix} \xi & \\ & 1 \end{bmatrix} g \right), \quad \text{where} \\
\widehat{\check{\Theta}'_{\check{\varphi}}(\check{F}')} (g) &:= \frac{1}{2} \int_{PB_{\mathbb{Q}}^{\times} \backslash PB_{\mathbb{A}}^{\times}} \int_{B_{\mathbb{A}}^{(\delta)}} \omega(g, \rho(\dot{\beta}\ddot{\beta}, \ddot{\beta})) \check{\varphi}(1) \\
&\quad \cdot \dot{\Psi}^{\eta}(\dot{\beta}\ddot{\beta}) \overline{\ddot{\Psi}^{\eta}(\ddot{\beta})} d^{(1)}\dot{\beta} d^{\times}\ddot{\beta} \\
&= \frac{1}{2} \int_{PB_{\mathbb{Q}}^{\times} \backslash PB_{\mathbb{A}}^{\times}} \int_{B_{\mathbb{A}}^{(\delta)}} \omega(g, \rho(\ddot{\beta}\dot{\beta}, \ddot{\beta})) \check{\varphi}(1) \\
&\quad \cdot \dot{\Psi}^{\eta}(\ddot{\beta}\dot{\beta}) \overline{\ddot{\Psi}^{\eta}(\ddot{\beta})} d^{(1)}\dot{\beta} d^{\times}\ddot{\beta} \\
&= \frac{1}{2} \int_{PB_{\mathbb{Q}}^{\times} \backslash PB_{\mathbb{A}}^{\times}} \tilde{\Psi}^{\eta}(g, \beta) \overline{\ddot{\Psi}^{\eta}(\beta)} d^{\times}\beta, \\
\tilde{\Psi}^{\eta}(g, \beta) &:= \int_{B_{\mathbb{A}}^{(\delta)}} \omega(g, \rho(\alpha, 1)) \check{\varphi}(1) \dot{\Psi}^{\eta}(\beta\alpha) d^{(1)}\alpha.
\end{aligned}$$

At this point our proof diverges from [34]. Using the calculations in §2.3, we

realize the last integral as the action of Hecke operators on $\dot{\Psi}^\eta$ and show

$$\tilde{\Psi}^\eta(g, \beta) = \widehat{F}(g) \dot{\Psi}^\eta(\beta),$$

where F is the unique Hecke normalized $\mathcal{H}^\star$ eigenform

$$\begin{aligned} F &\in \tilde{S}_{|k|}(1, (1, d_B N), \tilde{\chi}, N^\flat) \cap \pi^{\text{JL}}(\dot{\pi}) \quad \text{s.t.} \\ R_p(1, d_B N) F &= \dot{\varepsilon}_p F \quad \text{for all } p \mid N^\flat, \end{aligned}$$

and so

$$\check{\Theta}'_\varphi(\check{F}') = \langle \dot{\Psi}^\eta, \ddot{\Psi}^\eta \rangle F = \langle \dot{\Psi}, \ddot{\Psi} \rangle F.$$

It is clear from the definition of ω that

$$\tilde{\Psi}^\eta \left(\begin{bmatrix} 1 & x \\ & 1 \end{bmatrix} g, \beta \right) = \mathbf{e}_\mathbb{A}(x) \tilde{\Psi}^\eta(g, \beta) \quad \text{for } x \in \mathbb{A},$$

and since $\omega(z, \rho(z, 1)) = 1$,

$$\tilde{\Psi}^\eta(zg, \beta) = \tilde{\chi}(z) \tilde{\Psi}^\eta(g, \beta) \quad \text{for } z \in \mathbb{A}^\times.$$

Furthermore, the right K_v -types of $\tilde{\Psi}^\eta(g, \beta)$ are determined by Lemma 3 in §2.3. The remainder of our calculation consists identifying the Whittaker functions from §1.5 (with $M = d_B N$, $M^\flat = N^\flat$, $M^\sharp = d_B N^\sharp$, $M^\times = N^\times$).

If $p \nmid d_B N$, it suffices to consider $g = \begin{bmatrix} a & \\ & 1 \end{bmatrix}$:

$$\omega(g, \rho(\alpha, 1)) \check{\varphi}_p(1) = |a|_p \check{\varphi}_p(\alpha^\iota) = |a|_p \varphi_p(\alpha),$$

$$\begin{aligned} \tilde{\Psi}_p^\eta(g, \beta) &= |a|_p \int_{B_p^{(a)}} \varphi_p(\alpha) \dot{\Psi}_p^\eta(\beta\alpha) d_p^{(1)}\alpha \\ &= |a|_p \int_{B_p^{[a]}} \varphi_p(\alpha) \dot{\Psi}_p^\eta(\beta\alpha) d_p^\times\alpha \\ &= |a|_p T_p^{[a]} \dot{\Psi}_p^\eta(\beta) \\ &= |a|_p^{\frac{1}{2}} \left(\frac{|ap|_p^{\check{s}_p} - |ap|_p^{\check{s}_p}}{|p|_p^{\check{s}_p} - |p|_p^{\check{s}_p}} \right) \mathbb{I}_{\mathbb{Z}_p}(a) \dot{\Psi}_p^\eta(\beta) \\ &= W_p^0(g) \dot{\Psi}_p^\eta(\beta). \end{aligned}$$

If $p \mid N^\flat$, we consider g as before and also $g' = \begin{bmatrix} & a \\ -1 & \end{bmatrix}$:

$$\omega(g, \rho(\alpha, 1)) \check{\varphi}_p(1) = |a|_p \check{\varphi}_p(\alpha^\iota) = |a|_p \varphi_p(\alpha),$$

$$\begin{aligned}
\tilde{\Psi}_p^\eta(g, \beta) &= |a|_p T_p^{[a]} \dot{\Psi}_p^\eta(\beta) \\
&= |a|_p^{\frac{1}{2}} \left(\frac{|ap|_p^{\dot{s}_p} - |ap|_p^{\ddot{s}_p}}{|p|_p^{\dot{s}_p} - |p|_p^{\ddot{s}_p}} \right) \mathbb{I}_{\mathbb{Z}_p}(a) \dot{\Psi}_p^\eta(\beta) \\
&\quad + |a|_p^{\frac{1}{2}} \left(\frac{|a|_p^{\dot{s}_p} - |a|_p^{\ddot{s}_p}}{|p|_p^{\dot{s}_p} - |p|_p^{\ddot{s}_p}} \right) \mathbb{I}_{\mathbb{Z}_p}\left(\frac{a}{p}\right) R_p(\vec{N}) \dot{\Psi}_p^\eta(\beta) \\
&= (\dot{W}_p^0(g) + \overline{\varepsilon}_p \ddot{W}_p^0(g)) \dot{\Psi}_p^\eta(\beta),
\end{aligned}$$

$$\omega(g', \rho(\alpha, 1)) \check{\varphi}_p(1) = |a|_p \mathcal{F} \check{\varphi}_p(\alpha^\iota) = |a|_p \mathcal{F} \varphi_p(\alpha),$$

$$\begin{aligned}
\tilde{\Psi}_p^\eta(g', \beta) &= |a|_p \int_{B_p^{[a]}} \mathcal{F} \varphi_p(\alpha) \dot{\Psi}_p^\eta(\beta \alpha) d_p^\times \alpha \\
&= |ap|_p R_p(\vec{N})^{-1} T_p^{[ap]} \dot{\Psi}_p^\eta(\beta) \\
&= |ap|_p^{\frac{1}{2}} \left(\frac{|ap^2|_p^{\dot{s}_p} - |ap^2|_p^{\ddot{s}_p}}{|p|_p^{\dot{s}_p} - |p|_p^{\ddot{s}_p}} \right) \mathbb{I}_{\mathbb{Z}_p}(ap) R_p(\vec{N})^{-1} \dot{\Psi}_p^\eta(\beta) \\
&\quad + |a|_p^{\frac{1}{2}} \left(\frac{|ap|_p^{\dot{s}_p} - |ap|_p^{\ddot{s}_p}}{|p|_p^{\dot{s}_p} - |p|_p^{\ddot{s}_p}} \right) \mathbb{I}_{\mathbb{Z}_p}(a) \dot{\Psi}_p^\eta(\beta) \\
&= (\dot{W}_p^0(g') + \overline{\varepsilon}_p \ddot{W}_p^0(g')) \dot{\Psi}_p^\eta(\beta).
\end{aligned}$$

If $p \mid N^\sharp$,

$$\omega(g, \rho(\alpha, 1)) \check{\varphi}_p(1) = |a|_p \check{\varphi}_p(\alpha^\iota) = |a|_p \varphi_p(\alpha),$$

$$\begin{aligned}
\tilde{\Psi}_p^\eta(g, \beta) &= |a|_p T_p^{[a]} \dot{\Psi}_p^\eta(\beta) \\
&= |a|_p^{1+it_p} \mathbb{I}_{\mathbb{Z}_p}(a) \dot{\Psi}_p^\eta(\beta) \\
&= W_p^\sharp(g) \dot{\Psi}_p^\eta(\beta),
\end{aligned}$$

$$\omega(g', \rho(\alpha, 1)) \check{\varphi}_p(1) = |a|_p \mathcal{F} \check{\varphi}_p(\alpha^\iota) = |a|_p \mathcal{F} \varphi_p(\alpha),$$

$$\begin{aligned}
\tilde{\Psi}_p^\eta(g', \beta) &= |ap|_p R_p(\vec{N})^{-1} T_p^{[ap]} \dot{\Psi}_p^\eta(\beta) \\
&= -|a|_p^{it_p} |ap|_p \mathbb{I}_{\mathbb{Z}_p}(ap) \dot{\Psi}_p^\eta(\beta) \\
&= W_p^\sharp(g') \dot{\Psi}_p^\eta(\beta).
\end{aligned}$$

If $p \mid N^\chi$,

$$\omega(g, \rho(\alpha, 1)) \check{\varphi}_p(1) = |a|_p \check{\varphi}_p(\alpha^\iota) = |a|_p \varphi_p(\iota^{1-\eta_p} \alpha),$$

$$\begin{aligned}
\tilde{\Psi}_p^\eta(g, \beta) &= |a|_p \tilde{\chi}_p(a) \mathbb{I}_{\mathbb{Z}_p}(a) (T_p^\chi(a, 1) \dot{\Psi}_p^\eta)^{\eta_p}(\beta) \\
&= \tilde{\chi}_p(a) |a|_p^{\frac{1}{2}+it_p} \mathbb{I}_{\mathbb{Z}_p}(a) \dot{\Psi}_p^\eta(\beta) \\
&= W_p^\chi(g) \dot{\Psi}_p^\eta(\beta),
\end{aligned}$$

$$\begin{aligned}
\omega(g', \rho(\alpha, 1))\check{\varphi}_p(1) &= |a|_p \mathcal{F}\check{\varphi}_p(\alpha^\iota) = |a|_p \mathcal{F}\varphi_p(\iota^{1-\eta_p}\alpha), \\
\tilde{\Psi}_p^\eta(g', \beta) &= |ap^2|_p \overline{\mathcal{G}}_p \tilde{\chi}_p(-1) \mathbb{I}_{\mathbb{Z}_p}(ap) (T_p^\chi(p^{-1}, ap) \dot{\Psi}_p)^\eta(\beta), \\
&= \overline{\mathcal{G}}_p \tilde{\chi}_p(-1) |ap^2|_p^{\frac{1}{2}-it_p} \mathbb{I}_{\mathbb{Z}_p}(ap) \dot{\Psi}_p^\eta(\beta) \\
&= W_p^\chi(g') \dot{\Psi}_p^\eta(\beta).
\end{aligned}$$

If $p \mid d_B$,

$$\begin{aligned}
\omega(g, \rho(\alpha, 1))\check{\varphi}_p(1) &= |a|_p \check{\varphi}_p(\alpha^\iota) = |a|_p \varphi_p(\alpha), \\
\tilde{\Psi}_p^\eta(g, \beta) &= |a|_p T_p^{[a]} \dot{\Psi}_p^\eta(\beta) \\
&= |a|_p^{1+it_p} \mathbb{I}_{\mathbb{Z}_p}(a) \dot{\Psi}_p^\eta(\beta) \\
&= W_p^\sharp(g) \dot{\Psi}_p^\eta(\beta),
\end{aligned}$$

$$\begin{aligned}
\omega(g', \rho(\alpha, 1))\check{\varphi}_p(1) &= -|a|_p \mathcal{F}\check{\varphi}_p(\alpha^\iota) = -|a|_p \mathcal{F}\varphi_p(\alpha), \\
\tilde{\Psi}_p^\eta(g', \beta) &= -|ap|_p T_p(\varpi)^{-1} T_p^{[ap]} \dot{\Psi}_p^\eta(\beta) \\
&= -|a|_p^{it_p} |ap|_p \mathbb{I}_{\mathbb{Z}_p}(ap) \dot{\Psi}_p^\eta(\beta) \\
&= W_p^\sharp(g') \dot{\Psi}_p^\eta(\beta).
\end{aligned}$$

If $v = \infty$, we consider $g^\pm = \begin{bmatrix} \pm a & \\ & 1 \end{bmatrix}$, $a \in \mathbb{R}^+$:

$$\begin{aligned}
\omega(g^\pm, \rho(\alpha, 1))\check{\varphi}_\infty(1) &= a \check{\varphi}_\infty(\alpha^\iota), \\
\tilde{\Psi}_\infty^\eta(g^+, \beta) &= \int_{B_\infty^{(a)}} a^{1+\frac{|k|}{2}} \phi_{X_\infty}^{-\check{k}, a}(\alpha^\iota) \Psi_\infty^\eta(\beta\alpha) d_\infty^{(1)}\alpha \\
&= \int_{\mathbb{R}^+ \setminus B_\infty^+} a^{1+\frac{|k|}{2}} \phi_{X_\infty}^{\check{k}, a}(\alpha) \Psi_\infty^\eta(\beta\alpha) \frac{1}{2} d_\infty^\times \alpha \\
&= \frac{1}{2} a^{1+\frac{|k|}{2}} T_{X_\infty}^{\check{k}, a} \dot{\Psi}_\infty^\eta(\beta) \\
&= W_\infty^{|k|}(g^+) \dot{\Psi}_\infty^\eta(\beta), \\
\tilde{\Psi}_\infty^\eta(g^-, \beta) &= \frac{1}{2} a^{1+\frac{|k|}{2}} T_\infty^- T_{Y_\infty}^{\check{k}, a} \Psi_\infty^\eta(\beta) \\
&= W_\infty^{|k|}(g^-) \dot{\Psi}_\infty^\eta(\beta).
\end{aligned}$$

2.2.2 Adjoint of Shimizu's Lift

Let $\Psi = \dot{\Psi} = \ddot{\Psi}$, F be as in §2.2, and for $h' \in H_\mathbb{A}'^{(\delta)}$ define

$$\Theta_\varphi(\overline{F})(h') = \int_{G_\mathbb{Q}^{(1)} \setminus G_\mathbb{A}^{(\delta)}} \sum_{\alpha \in B_\mathbb{Q}} \omega(g, h') \varphi(\alpha) \overline{F(g)} dg.$$

Theorem 1.

$$\begin{aligned}\Theta_\varphi(\overline{F}) &= \frac{\|F\|^2}{\|\Psi\|^2} \overline{F'} \in L_0^2(H'_\mathbb{Q} \backslash H'_\mathbb{A}, \tilde{\chi}), \\ F'(\rho(\dot{\beta}, \ddot{\beta})\iota^\eta) &= \Psi^\eta(\dot{\beta}) \overline{\Psi^\eta(\ddot{\beta})}.\end{aligned}$$

Proof. Since the theta kernel is smooth, automorphic, and has moderate growth, it follows from Lemma 3 that

$$\begin{aligned}\Theta_\varphi(\overline{F})(\rho(\dot{\beta}, \ddot{\beta})\iota^\eta) &= \Theta_\varphi(\overline{F})(\rho(\dot{\beta}, \ddot{\beta})) \\ &\in \overline{\tilde{A}_k^\eta(d_B, \vec{N}, \tilde{\chi})} \times \tilde{A}_k^\eta(d_B, \vec{N}, \tilde{\chi}).\end{aligned}$$

Furthermore by Lemma 4, for $p \nmid d_B N$,

$$\Theta_\varphi(T_p^{[p]} \overline{F}) = \dot{T}_p^{[p]} \Theta_\varphi(\overline{F}) = (\ddot{T}_p^{[p]})^\vee \Theta_\varphi(\overline{F}),$$

and hence by strong multiplicity-one on $B_\mathbb{A}^\times$,

$$\Theta_\varphi(\overline{F}) \circ \rho \in \overline{\pi} \times \overline{\pi}^\vee = \overline{\pi} \times \pi.$$

Thus by Lemma 1, $\Theta_\varphi(\overline{F})$ is determined up to scaling. We compute its normalization using the adjoint identity below (compare with the similitude see-saw identity in [9]). $\square$

Lemma 2.

$$\left\langle \check{\Theta}'_\varphi(\check{F}'), F \right\rangle_{PG} = \left\langle \check{F}', \overline{\Theta_\varphi(\overline{F})} \right\rangle_{P\check{H}'}.$$

Proof.

$$\begin{aligned}\text{LHS} &= \frac{1}{2} \int_{G_\mathbb{Q} \mathbb{A}^\times \backslash G_\mathbb{A}} \frac{1}{2} \int_{\check{H}'_\mathbb{Q}(1) \backslash \check{H}'_\mathbb{A}(\delta)} \vartheta(g, h', \check{\varphi}) \check{F}'(h') \overline{F(g)} d^{(1)} h' d^\times g \\ &= \frac{1}{2} \int_{G_\mathbb{Q} \mathbb{R}^+ \backslash G_\mathbb{A}} \frac{1}{2} \int_{\check{H}'_\mathbb{Q}(1) \backslash \check{H}'_\mathbb{A}(\delta)} \vartheta(g, h', \check{\varphi}) \check{F}'(h') \overline{F(g)} d^{(1)} h' d^\times g \\ &= \int_{\mathbb{Q} \times \mathbb{R}^+ \backslash \mathbb{A}^\times} \int_{G_\mathbb{Q}^{(1)} \backslash G_\mathbb{A}^{(\delta)}} \frac{1}{2} \int_{\check{H}'_\mathbb{Q}(1) \backslash \check{H}'_\mathbb{A}(\delta)} \vartheta(g, h', \check{\varphi}) \check{F}'(h') \overline{F(g)} d^{(1)} h' d^{(1)} g d^\times \delta, \\ &= \text{RHS by symmetry.}\end{aligned}$$

$\square$

2.3 Local Data

With the notation of §2.2, define $\varphi, \check{\varphi} \in \mathcal{S}(B_{\mathbb{A}})$ by

$$\begin{aligned}\varphi_p(\beta) &= \frac{1}{\text{vol}^\times(\mathcal{O}_p^\times)} \mathbb{I}_{\mathcal{O}_p}(\beta) \begin{cases} 1 & \text{if } p \nmid N^\times, \\ \tilde{\chi}_p(\beta_{11}) \mathbb{I}_{\mathbb{Z}_p^\times}(\beta_{11}) & \text{if } p \mid N^\times, \end{cases} \\ \varphi_\infty(\beta) &= \frac{1}{\pi} X^k(\beta) \mathbf{e}(iP(\beta)), \\ \check{\varphi}_v(\beta) &= \omega(1, \iota_v^{\eta_v}) \varphi_v(\beta) = \varphi_v(\iota_v^{\eta_v} \beta), \quad \text{i.e.} \\ \check{\varphi}_p(\beta) &= \frac{1}{\text{vol}^\times(\mathcal{O}_p^\times)} \mathbb{I}_{\mathcal{O}_p}(\beta) \begin{cases} 1 & \text{if } p \nmid N^\times, \\ \tilde{\chi}_p(\beta_{11}) \mathbb{I}_{\mathbb{Z}_p^\times}(\beta_{11}) & \text{if } p \mid N^\times, \eta_p = 0, \\ \tilde{\chi}_p(\beta_{22}) \mathbb{I}_{\mathbb{Z}_p^\times}(\beta_{22}) & \text{if } p \mid N^\times, \eta_p = 1, \end{cases} \\ \check{\varphi}_\infty(\beta) &= \frac{1}{\pi} X^{\check{k}}(\beta) \mathbf{e}(iP(\beta)), \quad \check{k} = (-1)^{\eta_\infty} k.\end{aligned}$$

Now we compute the Fourier transforms of these functions in each case, and relate them to Hecke operators for $B_v^\times$.

If $p \nmid d_B N^\times$, writing $\beta = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$, $\alpha_{\vec{N}} = \begin{bmatrix} \dot{N} \\ \ddot{N} \end{bmatrix}$,

$$\begin{aligned}\check{\varphi}_p(\beta) &= \frac{1}{\text{vol}^\times(\mathcal{O}_p^\times)} \mathbb{I}_{\mathbb{Z}_p}(a) \mathbb{I}_{\dot{N}\mathbb{Z}_p}(b) \mathbb{I}_{\ddot{N}\mathbb{Z}_p}(c) \mathbb{I}_{\mathbb{Z}_p}(d), \\ \check{\varphi}_p|_{B_p^\times} &= \sum_{n \geq 0} T_p^{[p^n]}, \\ \mathcal{F}\check{\varphi}_p(\beta) &= \frac{1}{\text{vol}^\times(\mathcal{O}_p^\times)} \mathbb{I}_{\mathbb{Z}_p}(d) |\dot{N}|_p \mathbb{I}_{\frac{1}{\dot{N}}\mathbb{Z}_p}(c) |\ddot{N}|_p \mathbb{I}_{\frac{1}{\ddot{N}}\mathbb{Z}_p}(b) \mathbb{I}_{\mathbb{Z}_p}(a), \\ &= \frac{1}{\text{vol}^\times(\mathcal{O}_p^\times)} |N|_p \mathbb{I}_{\mathbb{Z}_p}(a) \mathbb{I}_{\mathbb{Z}_p}(\ddot{N}b) \mathbb{I}_{\mathbb{Z}_p}(\dot{N}c) \mathbb{I}_{\mathbb{Z}_p}(d), \\ &= \begin{cases} \check{\varphi}_p(\beta) & \text{if } p \nmid d_B N, \\ \frac{1}{p} \check{\varphi}_p(\alpha_{\vec{N}} \beta) & \text{if } p \mid N^\flat N^\sharp, \end{cases} \\ \mathcal{F}\check{\varphi}_p|_{B_p^\times} &= \begin{cases} \check{\varphi}_p|_{B_p^\times} & \text{if } p \nmid d_B N, \\ \frac{1}{p} R_p(\vec{N})^{-1} \check{\varphi}_p|_{B_p^\times} & \text{if } p \mid N^\flat N^\sharp. \end{cases}\end{aligned}$$

If $p \mid N^\times$, (note ι_p and $\mathcal{F}$ commute)

$$\begin{aligned}\varphi_p(\beta) &= \frac{1}{\text{vol}^\times(\mathcal{O}_p^\times)} \tilde{\chi}_p(a) \mathbb{I}_{\mathbb{Z}_p^\times}(a) \mathbb{I}_{\dot{N}\mathbb{Z}_p}(b) \mathbb{I}_{\ddot{N}\mathbb{Z}_p}(c) \mathbb{I}_{\mathbb{Z}_p}(d), \\ \varphi_p|_{B_p^\times} &= \sum_{n \geq 0} (\tilde{\chi}_p \circ \nu) \cdot \phi_{T_p(1, p^n)}^{\chi^{2,2}}, \\ \mathcal{F}\varphi_p(\beta) &= \frac{1}{\text{vol}^\times(\mathcal{O}_p^\times)} \frac{1}{p^2} \overline{\mathcal{G}_p \tilde{\chi}_p(-d)} \mathbb{I}_{\frac{1}{p}\mathbb{Z}_p^\times}(d) \mathbb{I}_{\frac{1}{p}\dot{N}\mathbb{Z}_p}(c) \mathbb{I}_{\frac{1}{p}\ddot{N}\mathbb{Z}_p}(b) \mathbb{I}_{\mathbb{Z}_p}(a), \\ \mathcal{F}\varphi_p|_{B_p^\times} &= \frac{1}{p^2} \overline{\mathcal{G}_p \tilde{\chi}_p(-1)} \sum_{n \geq 0} \phi_{T_p(p^n, p^{-1})}^{\chi^{2,2}}.\end{aligned}$$

If $p \mid d_B$, $R_p^* := \{\alpha \in D_p; \forall \beta \in R_p, \text{tr}(\alpha\beta) \in \mathbb{Z}_p\} = \varpi_p^{-1} R_p$,

$$\begin{aligned}\check{\varphi}_p &= \frac{1}{\text{vol}^\times(R_p)} \mathbb{I}_{R_p}, \\ \check{\varphi}_p|_{B_p^\times} &= \sum_{n \geq 0} T_p^{[p^n]}, \\ -\omega(J)\check{\varphi}_p &= \mathcal{F}\check{\varphi}_p = \frac{1}{\text{vol}^\times(R_p)} \frac{1}{p} \mathbb{I}_{(R_p^*)^\iota} = \frac{1}{\text{vol}^\times(R_p)} \frac{1}{p} \mathbb{I}_{\varpi_p^{-1} R_p}, \\ \mathcal{F}\check{\varphi}_p|_{B_p^\times} &= \frac{1}{p} T_p(\varpi_p)^{-1} \check{\varphi}_p|_{B_p^\times}.\end{aligned}$$

If $v = \infty$,

$$\begin{aligned}\check{\varphi}_\infty|_{B_\infty^+} &= \nu^{\frac{|k|}{2}} \cdot \phi_{X_\infty}^{-k, \nu}, \\ T_\infty^- \check{\varphi}_\infty|_{B_\infty^-} &= \nu^{\frac{|k|}{2}} \cdot \phi_{Y_\infty}^{-k, \nu}.\end{aligned}$$

Then

$$\begin{aligned}D\omega\left(\begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix}\right) &= 2\pi i (X\overline{X} - Y\overline{Y}) \cdot, \\ -D\omega\left(\begin{bmatrix} 0 & 0 \\ 1 & 0 \end{bmatrix}\right) &= \frac{1}{2\pi i} \left(\frac{\partial}{\partial X} \frac{\partial}{\partial \overline{X}} - \frac{\partial}{\partial Y} \frac{\partial}{\partial \overline{Y}}\right), \\ D\omega(J) \check{\varphi}_\infty &= i|k| \check{\varphi}_\infty \\ \implies \omega(\kappa_\theta) \check{\varphi}_\infty &= e^{i|k|\theta} \check{\varphi}_\infty.\end{aligned}$$

Lemma 3. For $\kappa \in K_p(1, (1, d_B N))$ and $\dot{\kappa}, \ddot{\kappa} \in K_p(d_B, \vec{N})$ s.t. $\det \kappa = \nu(\dot{\kappa})/\nu(\ddot{\kappa})$, (Note $\tilde{\chi}_p$ has different definitions on these two groups.)

$$\begin{aligned}\omega(\kappa, \rho(\dot{\kappa}, \ddot{\kappa}))\check{\varphi}_p &= \tilde{\chi}_p(\kappa) \overline{\tilde{\chi}_p(\iota_p^{\eta_p} \dot{\kappa})} \tilde{\chi}_p(\iota_p^{\eta_p} \ddot{\kappa}) \check{\varphi}_p, \\ \omega(\kappa_\theta, \rho(\kappa_\theta, \kappa_{\bar{\theta}}))\check{\varphi}_\infty &= e^{i(|k|\theta - \check{k}\bar{\theta} + \check{k}\bar{\theta})} \check{\varphi}_\infty.\end{aligned}$$

Proof. First consider diagonal $\kappa \in K_p(1, (1, d_B N))$:

$$\begin{aligned}\omega(\kappa, \rho(\dot{\kappa}, \ddot{\kappa}))\check{\varphi}_p(\beta) &= \check{\varphi}_p(\kappa_{11} \dot{\kappa}^{-1} \beta \ddot{\kappa}) \\ &= \tilde{\chi}_p(\kappa) \overline{\tilde{\chi}_p(\iota_p^{\eta_p} \dot{\kappa})} \tilde{\chi}_p(\iota_p^{\eta_p} \ddot{\kappa}) \check{\varphi}_p.\end{aligned}$$

Then it suffices to check

$$\begin{aligned}\omega\left(\begin{bmatrix} 1 & x \\ & 1 \end{bmatrix}\right)\check{\varphi}_p &= \check{\varphi}_p \quad \text{for } x \in \mathbb{Z}_p, \\ \omega\left(\begin{bmatrix} 1 & x \\ & 1 \end{bmatrix}\right)\mathcal{F}\check{\varphi}_p &= \mathcal{F}\check{\varphi}_p \quad \text{for } x \in d_B N \mathbb{Z}_p.\end{aligned}$$

The archimedean statement follows from previous calculations. □

Lemma 4. For $p \nmid d_B N$, $\alpha = \begin{bmatrix} 1 & \\ & p \end{bmatrix}$,

$$\begin{aligned} \int_{\mathrm{SL}_2(\mathbb{Z}_p)} \omega(\alpha \kappa \alpha^{-1}) \varphi_p(\beta) d_p^{(1)} \kappa &= \int_{K_p^{(1)}(d_B, \vec{N})} \omega(\alpha, \rho(\dot{\kappa}^{-1} \alpha, 1)) \varphi_p(\beta) d_p^{(1)} \dot{\kappa} \\ &= \int_{K_p^{(1)}(d_B, \vec{N})} \omega(\alpha, \rho(1, \ddot{\kappa} \alpha^{-1})) \varphi_p(\beta) d_p^{(1)} \ddot{\kappa}. \end{aligned}$$

Proof. Since $\omega(\alpha \kappa \alpha^{-1}) \varphi_p = \varphi_p$ for $\kappa \in K_p^{(1)}(1, (p, \frac{1}{p}))$, the κ integral reduces to an average over the $K_p^{(1)}(1, (1, 1))/K_p^{(1)}(1, (p, 1))$ coset representatives,

$$\begin{bmatrix} 1 & x \\ & 1 \end{bmatrix} \quad (x \bmod p), \quad \begin{bmatrix} & 1 \\ -1 & \end{bmatrix},$$

and hence evaluates to

$$\varphi_p'(\beta) = \frac{1}{p+1} (p \mathbb{I}_{\mathcal{O}_p^{(p\mathbb{Z}_p)}(d_B, \vec{N})}(\beta) + p^2 \mathbb{I}_{p\mathcal{O}_p(d_B, \vec{N})}(\beta)).$$

Now the $\dot{\kappa}$ and $\ddot{\kappa}$ integrals evaluate to

$$\begin{aligned} \int_{K_p^{(1)}(d_B, \vec{N})} p \varphi_p(\alpha^{-1} \dot{\kappa} \beta) d_p^{(1)} \dot{\kappa} &= \frac{p}{p+1} (T_p^{[p]} * \mathbb{I}_{\mathcal{O}_p})(\beta), \\ \int_{K_p^{(1)}(d_B, \vec{N})} p \varphi_p(\beta \ddot{\kappa} \alpha^{-1}) d_p^{(1)} \ddot{\kappa} &= \frac{p}{p+1} (\mathbb{I}_{\mathcal{O}_p} * T_p^{[p]})(\beta). \end{aligned}$$

It follows from the calculations in §1.3.2 (extending by continuity to B_v) that

$$\begin{aligned} \mathbb{I}_{\mathcal{O}_p^\times} &= \mathbb{I}_{\mathcal{O}_p} * (T_p(1) - T_p^{[p]} + p T_p(p)), \\ \mathbb{I}_{\mathcal{O}_p} * T_p^{[p]} &= \mathbb{I}_{\mathcal{O}_p} - \mathbb{I}_{\mathcal{O}_p^\times} + p \mathbb{I}_{p\mathcal{O}_p} = T_p^{[p]} * \mathbb{I}_{\mathcal{O}_p} \\ &= \mathbb{I}_{\mathcal{O}_p^{(p\mathbb{Z}_p)}} + p \mathbb{I}_{p\mathcal{O}_p} = \frac{p+1}{p} \varphi_p'. \end{aligned}$$

□

Chapter 3

L-functions

In this chapter we compute special values of adjoint and triple product L-functions. First we describe the relevant Langlands parameters (admissible representations of Weil or Weil-Deligne groups), which determine canonical L and ε factors. Then we compute special values of the global L -functions using the Rankin–Selberg and Garrett/Piatetski-Shapiro–Rallis integral representations.

3.1 Local Langlands Correspondence

Let W_F denote the Weil group of the local field F (as in [39, 22, 32]) and r_F the local class-field theory isomorphism $r_F : F^\times \xrightarrow{\sim} W_F^{\text{ab}}$. If χ is a character of $F^\times$, denote by χ^r the corresponding character of W_F . In particular, write $\|\cdot\| = \|\cdot\|_F^r$.

Recall that $W_{\mathbb{C}} = \mathbb{C}^\times$ is abelian and $r_{\mathbb{C}}$ is the identity (so $\|z\| = |z|_{\mathbb{C}}$), and

$$W_{\mathbb{R}} = \mathbb{C}^\times \cup j\mathbb{C}^\times, \quad j^2 = -1, \quad jzj^{-1} = \bar{z},$$

has closed commutator subgroup $W_{\mathbb{R}}^c = \mathbb{C}^{(1)}$, with $r_{\mathbb{R}}$ being defined by

$$\mathbb{R}^+ \ni r \mapsto r^{\frac{1}{2}} \mathbb{C}^{(1)}, \quad -1 \mapsto j \mathbb{C}^{(1)} \quad (\text{so } \|z\| = \|jz\| = |z|_{\mathbb{C}}).$$

As described in [21] (with a normalization error that we have corrected), all one-dimensional representations of $W_{\mathbb{R}}$ are isomorphic to some $\varrho^1 = (s, \delta)_{\mathbb{R}}^1$,

$$\varrho^1(z) = \|z\|^s, \quad \varrho^1(j) = (-1)^\delta, \quad \text{for } s \in \mathbb{C}, \delta \in \{0, 1\},$$

and all irreducible two-dimensional semi-simple representations to some $\varrho^2 = (s, l)_{\mathbb{R}}^2$,

$$\varrho^2(re^{i\theta}) = \begin{bmatrix} r^{2s}e^{il\theta} & \\ & r^{2s}e^{-il\theta} \end{bmatrix}, \quad \varrho^2(j) = \begin{bmatrix} & (-1)^l \\ 1 & \end{bmatrix}, \quad \text{for } s \in \mathbb{C}, l \in \mathbb{Z}^+.$$

If we allow $l \in \mathbb{Z}$, then

$$\begin{aligned} (s, 0)_{\mathbb{R}}^2 &\simeq (s, 0)_{\mathbb{R}}^1 \oplus (s, 1)_{\mathbb{R}}^1, & (s, l)_{\mathbb{R}}^2 &\simeq (s, -l)_{\mathbb{R}}^2, \\ (s_1, \delta_1)_{\mathbb{R}}^1 \otimes (s_2, \delta_2)_{\mathbb{R}}^1 &\simeq (s_1 + s_2, \delta)_{\mathbb{R}}^1, & \delta &\equiv \delta_1 + \delta_2 \pmod{2}, \\ (s_1, l_1)_{\mathbb{R}}^2 \otimes (s_2, l_2)_{\mathbb{R}}^2 &\simeq (s_1 + s_2, l_1 + l_2)_{\mathbb{R}}^2 \oplus (s_1 + s_2, l_1 - l_2)_{\mathbb{R}}^2, \\ (s_1, \delta_1)_{\mathbb{R}}^1 \otimes (s_2, l_2)_{\mathbb{R}}^2 &\simeq (s_1 + s_2, l_2)_{\mathbb{R}}^2. \end{aligned}$$

The standard local factors are defined as follows:

$$\begin{aligned} \zeta_{\mathbb{R}}(s) &= \pi^{-\frac{s}{2}} \Gamma(\frac{s}{2}), & \zeta_{\mathbb{C}}(s) &= \zeta_{\mathbb{R}}(s) \zeta_{\mathbb{R}}(s+1) \\ & & &= (2\pi)^{-s} \Gamma(s), \\ L(s, \varrho^1) &= \zeta_{\mathbb{R}}(s + s_{\infty} + \delta_{\infty}), & L(s, \varrho^2) &= \zeta_{\mathbb{C}}(s + s_{\infty} + \frac{l_{\infty}}{2}), \\ C(t, \varrho^1) &= (1 + |it + s_{\infty}|), & C(t, \varrho^2) &= (1 + |it + s_{\infty} + \frac{l_{\infty}}{2}|)^2, \\ \varepsilon(s, \varrho^1, e_{\infty}) &= i^{\delta_{\infty}}, & \varepsilon(s, \varrho^2, e_{\infty}) &= i^{l_{\infty}+1}. \end{aligned}$$

Now let $W'_{\mathbb{Q}_p}$ denote the Weil-Deligne group of $\mathbb{Q}_p$. We implicitly identify the characters of $W_{\mathbb{Q}_p}$ and $W'_{\mathbb{Q}_p}$. Recall from [39, 22] the indecomposable admissible representation $\mathbf{sp}^n$ of $W'_{\mathbb{Q}_p}$ on $\mathbb{C}\{e_0, \dots, e_{n-1}\}$ defined by

$$\mathbf{sp}^n(w) e_i = \|w\|^i e_i, \quad \mathbf{sp}^n(N) e_i = \begin{cases} e_{i+1} & \text{if } i < n-1, \\ 0 & \text{if } i = n-1. \end{cases}$$

Note $(\mathbf{sp}^n)^{\vee} \simeq \|\cdot\|^{1-n} \otimes \mathbf{sp}^n$. Since $\mathbf{sp}^n \simeq \text{Sym}^{n-1} \mathbf{sp}^2$, it follows that

$$\begin{aligned} \mathbf{sp}^m \otimes \mathbf{sp}^n &\simeq \bigoplus_{i=0}^{n-1} \|\cdot\|^i \otimes \mathbf{sp}^{m+n-2i-1} \quad \forall m \geq n, \\ \otimes^2 \mathbf{sp}^2 &\simeq \|\cdot\|^1 \oplus \mathbf{sp}^3, \\ \otimes^3 \mathbf{sp}^2 &\simeq \oplus^2 (\|\cdot\|^1 \otimes \mathbf{sp}^2) \oplus \mathbf{sp}^4. \end{aligned}$$

Also consider a character $\tilde{\chi}_p$ of $\mathbb{Q}_p^{\times}$ with conductor $p^k \mathbb{Z}_p$ for $k > 0$. The standard local factors we need are then defined as follows: $\zeta_p(s) = (1 - p^{-s})^{-1}$,

$$\begin{aligned} \varrho &= \|\cdot\|^{s_p} \otimes \mathbf{sp}^n, & \varrho' &= \tilde{\chi}_p^r, \\ L(s, \varrho) &= \zeta_p(s + s_p + n - 1), & L(s, \varrho') &= 1, \\ C(\varrho) &= p^{n-1}, & C(\varrho') &= p^k, \\ \varepsilon(s, \varrho, e_p) &= \prod_{i=0}^{n-2} \frac{\zeta_p(-s - s_p - i)}{\zeta_p(s + s_p + i)} & \varepsilon(s, \varrho', e_p) &= p^{-ks} \int_{p^{-k} \mathbb{Z}_p^{\times}} \tilde{\chi}_p^{-1}(x) e_p(x) dx. \\ &= \left(-p^{-s-s_p-\frac{n-2}{2}} \right)^{n-1}, \end{aligned}$$

3.1.1 Adjoint Parameters

Consider a representation π_v of $\mathrm{GL}_2(\mathbb{Q}_v)$ as in §1.4. Below we list the corresponding Langlands parameters ϱ and their local factors, plus those of compositions with the adjoint representation $\mathrm{Ad} : \mathrm{GL}_2(\mathbb{C}) \rightarrow \mathrm{GL}_3(\mathbb{C})$, $\mathrm{Ad}\varrho = \mathrm{Ad} \circ \varrho \simeq \varrho \otimes \varrho^\vee \oplus 1$.

Archimedean

If $k = 0$, then $\pi_\infty \simeq \pi(\mathrm{sgn}^{\delta_\infty} | \cdot |_\infty^{s_\infty}, \mathrm{sgn}^{\delta_\infty} | \cdot |_\infty^{-s_\infty})$, $\delta_\infty \in \{0, 1\}$, corresponds to

$$\begin{aligned} \varrho^0 &\simeq (s_\infty, \delta_\infty)_{\mathbb{R}}^1 \oplus (-s_\infty, \delta_\infty)_{\mathbb{R}}^1, \text{ with} \\ L(s, \varrho^0) &= \zeta_{\mathbb{R}}(s + s_\infty + \delta_\infty) \zeta_{\mathbb{R}}(s - s_\infty + \delta_\infty), \\ C(t, \varrho^0) &= (1 + |it + s_\infty|)(1 + |it - s_\infty|), \\ \varepsilon(s, \varrho^0, \mathrm{e}_\infty) &= (-1)^{\delta_\infty}, \text{ and} \\ \mathrm{Ad}\varrho^0 &\simeq (2s_\infty, 0)_{\mathbb{R}}^1 \oplus (0, 0)_{\mathbb{R}}^1 \oplus (-2s_\infty, 0)_{\mathbb{R}}^1, \text{ with} \\ L(s, \mathrm{Ad}\varrho^0) &= \zeta_{\mathbb{R}}(s + 2s_\infty) \zeta_{\mathbb{R}}(s) \zeta_{\mathbb{R}}(s - 2s_\infty), \\ C(t, \mathrm{Ad}\varrho^0) &= (1 + |it + 2s_\infty|)(1 + |it|)(1 + |it - 2s_\infty|), \\ \varepsilon(s, \mathrm{Ad}\varrho^0, \mathrm{e}_\infty) &= 1. \end{aligned}$$

If $k \geq 2$, then $\pi_\infty \simeq \sigma(\mathrm{sgn}^k | \cdot |_\infty^{s_\infty}, | \cdot |_\infty^{-s_\infty})$, $s_\infty = \frac{k-1}{2}$, corresponds to

$$\begin{aligned} \varrho^k &\simeq (0, k-1)_{\mathbb{R}}^2, \text{ with} \\ L(s, \varrho^k) &= \zeta_{\mathbb{C}}(s + s_\infty), \\ C(t, \varrho^k) &= (1 + |it + s_\infty|)^2, \\ \varepsilon(s, \varrho^k, \mathrm{e}_\infty) &= i^k, \text{ and} \\ \mathrm{Ad}\varrho^k &\simeq (0, 2k-2)_{\mathbb{R}}^2 \oplus (0, 1)_{\mathbb{R}}^1, \text{ with} \\ L(s, \mathrm{Ad}\varrho^k) &= \zeta_{\mathbb{C}}(s + 2s_\infty) \zeta_{\mathbb{R}}(s + 1), \\ C(t, \mathrm{Ad}\varrho^k) &= (1 + |it + 2s_\infty|)^2 (1 + |it|), \\ \varepsilon(s, \mathrm{Ad}\varrho^k, \mathrm{e}_\infty) &= (-1)^k. \end{aligned}$$

Non-Archimedean

If $p \nmid M^\sharp M^\chi$, then $\pi_p \simeq \pi(| \cdot |_p^{\dot{s}_p}, | \cdot |_p^{\ddot{s}_p})$ corresponds to

$$\begin{aligned} \varrho^0 &\simeq \| \cdot \|_p^{\dot{s}_p} \oplus \| \cdot \|_p^{\ddot{s}_p}, \text{ with} \\ L(s, \varrho^0) &= \zeta_p(s + \dot{s}_p) \zeta_p(s + \ddot{s}_p), \\ \varepsilon(s, \varrho^0, \mathrm{e}_p) &= 1, \\ C(\varrho^0) &= 1, \text{ and} \\ \mathrm{Ad}\varrho^0 &\simeq \| \cdot \|_p^{\dot{s}_p - \ddot{s}_p} \oplus \| \cdot \|_p^0 \oplus \| \cdot \|_p^{\ddot{s}_p - \dot{s}_p}, \text{ with} \\ L(s, \mathrm{Ad}\varrho^0) &= \zeta_p(s + \dot{s}_p - \ddot{s}_p) \zeta_p(s) \zeta_p(s + \ddot{s}_p - \dot{s}_p), \\ \varepsilon(s, \mathrm{Ad}\varrho^0, \mathrm{e}_p) &= 1, \\ C(\mathrm{Ad}\varrho^0) &= 1. \end{aligned}$$

If $p \mid M^\sharp$, then $\pi_p \simeq \sigma(|p|^{-\frac{1}{2}+it_p}, |p|^{\frac{1}{2}+it_p})$ corresponds to

$$\begin{aligned} \varrho^\sharp &\simeq \| \|^{-\frac{1}{2}+it_p} \otimes \mathfrak{sp}^2 & \text{and } \text{Ad}\varrho^\sharp &\simeq \| \|^{-1} \otimes \mathfrak{sp}^3, \text{ with} \\ L_p(s, \varrho^\sharp) &= \zeta_p(s + \tfrac{1}{2} + it_p), & L(s, \text{Ad}\varrho^\sharp) &= \zeta_p(s + 1), \\ \varepsilon_p(s, \varrho^\sharp, \mathbf{e}_p) &= (-p^{-it_p})p^{-s+\frac{1}{2}}, & \varepsilon(s, \text{Ad}\varrho^\sharp, \mathbf{e}_p) &= p^{-2s+1}, \\ C(\varrho^\sharp) &= p, & C(\text{Ad}\varrho^\sharp) &= p^2. \end{aligned}$$

If $p \mid M^\chi$, then $\pi_p \simeq \pi(\tilde{\chi}_p |p|^{\frac{it_p}{p}}, |p|^{-it_p})$ corresponds to

$$\begin{aligned} \varrho^\chi &\simeq \tilde{\chi}_p^r \| \|^{it_p} \oplus \| \|^{-it_p}, \text{ with} \\ L(s, \varrho^\chi) &= \zeta_p(s - it_p), \\ \varepsilon(s, \varrho^\chi, \mathbf{e}_p) &= \left(p^{-it_p-\frac{1}{2}} \int_{\frac{1}{p}\mathbb{Z}_p^\times} \overline{\tilde{\chi}_p(x)} \mathbf{e}_p(x) dx \right) p^{-s+\frac{1}{2}}, \\ C(\varrho^\chi) &= p, \text{ and} \\ \text{Ad}\varrho^\chi &\simeq \tilde{\chi}_p^r \| \|^{2it_p} \oplus \| \|^0 \oplus \overline{\tilde{\chi}_p^r} \| \|^{-2it_p}, \text{ with} \\ L(s, \text{Ad}\varrho^\chi) &= \zeta_p(s), \\ \varepsilon(s, \text{Ad}\varrho^\chi, \mathbf{e}_p) &= \tilde{\chi}_p(-1) p^{-2s+1}, \\ C(\text{Ad}\varrho^\chi) &= p^2. \end{aligned}$$

3.1.2 Triple Product Parameters

Archimedean

If $k_1 = k_2 = k_3 = 0$, let $\{0, 1\} \ni \delta \equiv \delta_1 + \delta_2 + \delta_3 \pmod{2}$, so that

$$\begin{aligned} \varrho &:= \varrho_1^0 \otimes \varrho_2^0 \otimes \varrho_3^0 \simeq \bigoplus_{\{\pm\}^3} (\pm s_1 \pm s_2 \pm s_3, \delta)_{\mathbb{R}}^1, \text{ with} \\ L(s, \varrho) &= \prod_{\{\pm\}^3} \zeta_{\mathbb{R}}(s \pm s_1 \pm s_2 \pm s_3 + \delta), \\ C(t, \varrho) &= \prod_{\{\pm\}^3} (1 + |it \pm s_1 \pm s_2 \pm s_3|), \\ \varepsilon(s, \varrho, \mathbf{e}_\infty) &= 1. \end{aligned}$$

If $k = |k_1| = |k_2| \geq 2$ and $k_3 = 0$,

$$\begin{aligned} \varrho &:= \varrho_1^k \otimes \varrho_2^k \otimes \varrho_3^0 \simeq (s_3, 2k-2)_{\mathbb{R}}^2 \oplus (s_3, 0)_{\mathbb{R}}^2 \\ &\quad \oplus (-s_3, 2k-2)_{\mathbb{R}}^2 \oplus (-s_3, 0)_{\mathbb{R}}^2, \text{ with} \\ L(s, \varrho) &= \zeta_{\mathbb{C}}(s + s_3 + k - 1) \zeta_{\mathbb{C}}(s + s_3) \\ &\quad \cdot \zeta_{\mathbb{C}}(s - s_3 + k - 1) \zeta_{\mathbb{C}}(s - s_3), \\ C(t, \varrho) &= (1 + |it + s_3 + k - 1|)^2 (1 + |it + s_3|)^2 \\ &\quad \cdot (1 + |it - s_3 + k - 1|)^2 (1 + |it - s_3|)^2, \\ \varepsilon(s, \varrho, \mathbf{e}_\infty) &= 1. \end{aligned}$$

If $|k_1| = |k_2| + |k_3| > |k_2| \geq |k_3| \geq 2$,

$$\begin{aligned}\varrho &:= \varrho_1^{|k_1|} \otimes \varrho_2^{|k_2|} \otimes \varrho_3^{|k_3|} \simeq (0, 2|k_1| - 3)_{\mathbb{R}}^2 \oplus (0, 2|k_2| - 1)_{\mathbb{R}}^2 \\ &\quad \oplus (0, 2|k_3| - 1)_{\mathbb{R}}^2 \oplus (0, 1)_{\mathbb{R}}^2, \text{ with} \\ L(s, \varrho) &= \zeta_{\mathbb{C}}(s + |k_1| - \tfrac{3}{2}) \zeta_{\mathbb{C}}(s + |k_2| - \tfrac{1}{2}) \\ &\quad \cdot \zeta_{\mathbb{C}}(s + |k_3| - \tfrac{1}{2}) \zeta_{\mathbb{C}}(s + \tfrac{1}{2}), \\ C(t, \varrho) &= (1 + |it + |k_1| - \tfrac{3}{2}|)^2 (1 + |it + |k_2| - \tfrac{1}{2}|)^2 \\ &\quad \cdot (1 + |it + |k_3| - \tfrac{1}{2}|)^2 (1 + |it + \tfrac{1}{2}|)^2, \\ \varepsilon(s, \varrho, e_{\infty}) &= 1.\end{aligned}$$

Non-Archimedean

$$\begin{aligned}\varrho_1^0 \otimes \varrho_2^0 &\simeq \bigoplus_{\eta \in \{\cdot, \dots\}^2} \|\| s_1^{\eta_1} + s_2^{\eta_2}, \\ \varrho_1^0 \otimes \varrho_2^0 \otimes \varrho_3^0 &\simeq \bigoplus_{\eta \in \{\cdot, \dots\}^3} \|\| s_1^{\eta_1} + s_2^{\eta_2} + s_3^{\eta_3}, \\ L_p(s, \varrho_1^0 \otimes \varrho_2^0 \otimes \varrho_3^0) &= \prod_{\eta \in \{\cdot, \dots\}^3} \zeta_p(s + s_1^{\eta_1} + s_2^{\eta_2} + s_3^{\eta_3}), \\ \varepsilon_p(s, \varrho_1^0 \otimes \varrho_2^0 \otimes \varrho_3^0, e_p) &= 1, \\ C(\varrho_1^0 \otimes \varrho_2^0 \otimes \varrho_3^0) &= 1, \\ \varrho_1^0 \otimes \varrho_2^0 \otimes \varrho_3^{\sharp} &\simeq \bigoplus_{\eta \in \{\cdot, \dots\}^2} \|\| s_1^{\eta_1} + s_2^{\eta_2} - \tfrac{1}{2} + it_3 \otimes \mathfrak{sp}^2, \\ \varrho_1^{\sharp} \otimes \varrho_2^{\sharp} &\simeq \|\|^{it_1 + it_2} \oplus (\|\|^{-1 + it_1 + it_2} \otimes \mathfrak{sp}^3), \\ \varrho_1^{\sharp} \otimes \varrho_2^{\sharp} \otimes \varrho_3^{\sharp} &\simeq \oplus^2 (\|\|^{-\frac{1}{2} + i(t_1 + t_2 + t_3)} \otimes \mathfrak{sp}^2) \oplus (\|\|^{-\frac{3}{2} + i(t_1 + t_2 + t_3)} \otimes \mathfrak{sp}^4), \\ L(s, \varrho_1^{\sharp} \otimes \varrho_2^{\sharp} \otimes \varrho_3^{\sharp}) &= \zeta_p(s + \tfrac{1}{2} + i(t_1 + t_2 + t_3))^2 \zeta_p(s + \tfrac{3}{2} + i(t_1 + t_2 + t_3)), \\ \varepsilon(s, \varrho_1^{\sharp} \otimes \varrho_2^{\sharp} \otimes \varrho_3^{\sharp}, e_p) &= (-p^{-5i(t_1 + t_2 + t_3)}) p^{-5s + \frac{5}{2}}, \\ C(\varrho_1^{\sharp} \otimes \varrho_2^{\sharp} \otimes \varrho_3^{\sharp}) &= p^5, \\ \varrho_1^{\sharp} \otimes \varrho_2^{\sharp} \otimes \varrho_3^0 &\simeq \|\|^{it_1 + it_2 + \dot{s}_3} \oplus (\|\|^{-1 + it_1 + it_2 + \dot{s}_3} \otimes \mathfrak{sp}^3) \\ &\quad \oplus \|\|^{it_1 + it_2 + \ddot{s}_3} \oplus (\|\|^{-1 + it_1 + it_2 + \ddot{s}_3} \otimes \mathfrak{sp}^3).\end{aligned}$$

3.2 Zeta Integrals

3.2.1 Rankin–Selberg

Lemma 5. *Let $F \in \tilde{S}_k(1, (1, M), \tilde{\chi}, M^b)$ be a Hecke normalized $\mathcal{H}^*$ eigenform, with Langlands parameters $\varrho = (\varrho_v)$. Then*

$$\langle F, F \rangle = \int_{\mathbb{A}^\times} \left| \hat{F} \left(\begin{bmatrix} a & \\ & 1 \end{bmatrix} \right) \right|^2 |a|_{\mathbb{A}}^{-1} da = \frac{\prod_v c_v}{\zeta^*(2)} L^*(1, \text{Ad} \varrho),$$

$$c_\infty = \begin{cases} 1 & \text{if } k = 0, \\ 2^{-|k|-1} & \text{if } k \neq 0, \end{cases} \quad c_p = \begin{cases} 1 & \text{if } p \nmid M, \\ \frac{p}{p+1} & \text{if } p \mid M^\sharp M^\chi, \\ 2 \left(1 + \frac{\overline{\varepsilon}_p \lambda_p}{p+1} \right) & \text{if } p \mid M^b. \end{cases}$$

In terms of the corresponding classical Hecke form $f \in S_k(1, (1, M), \chi, M^b)$,

$$\int_{\Gamma_0(M) \backslash \mathfrak{H}} |f(z)|^2 \frac{dx dy}{y^2} = 2c_\infty M \left(\prod_{p \mid M^b} 2 \left(1 + \frac{1 + \overline{\varepsilon}_p \lambda_p}{p} \right) \right) L^*(1, \text{Ad} \varrho).$$

Proof. Suppose $M^b = 1$ for now, and define the $PG_{\mathbb{A}}$ Eisenstein series

$$E(s, g, \xi) = \sum_{\gamma \in PT_{\mathbb{Q}} \backslash PG_{\mathbb{Q}}} \xi(s, \gamma g),$$

$$\xi_v \left(s, \begin{bmatrix} a & x \\ & 1 \end{bmatrix} \kappa \right) = |a|_v^s \frac{1}{\text{vol}_v} \mathbb{I}_{PK_v(1, (1, M))}(\kappa),$$

$$\kappa \in PK_v(1, (1, 1)), \quad \text{vol}_v = \text{vol}_v(PK_v(1, (1, M))).$$

Now consider the Rankin–Selberg zeta integral,

$$\begin{aligned} Z(s, F \times \overline{F}, \xi) &= \int_{PG_{\mathbb{Q}} \backslash PG_{\mathbb{A}}} |F(g)|^2 E(s, g, \xi) dg \\ &= \int_{PT_{\mathbb{Q}} \backslash PG_{\mathbb{A}}} |F(g)|^2 \xi(s, g) dg \\ &= \int_{\mathbb{Q}^\times \backslash \mathbb{A}^\times} \int_{\mathbb{Q} \backslash \mathbb{A}} \left| F \left(\begin{bmatrix} a & x \\ & 1 \end{bmatrix} \right) \right|^2 |a|_{\mathbb{A}}^{s-1} d^\times a dx \\ &= \int_{\mathbb{Q}^\times \backslash \mathbb{A}^\times} \sum_{\xi \in \mathbb{Q}^\times} \left| \hat{F} \left(\begin{bmatrix} \xi a & \\ & 1 \end{bmatrix} \right) \right|^2 |a|_{\mathbb{A}}^{s-1} d^\times a \\ &= \prod_v Z_v(s, \hat{F}_v \times \hat{\overline{F}}_v), \\ Z_v(s, \hat{F}_v \times \hat{\overline{F}}_v) &= \int_{\mathbb{Q}_v^\times} \left| \hat{F}_v \left(\begin{bmatrix} a & \\ & 1 \end{bmatrix} \right) \right|^2 |a|_v^{s-1} d_v^\times a. \end{aligned}$$

These local factors are easy to compute: If $k = 0$,

$$\begin{aligned} Z_\infty(s, W_p^0 \times \check{W}_p^0) &= \zeta_{\mathbb{R}}(s + 2s_\infty) \zeta_{\mathbb{R}}(s)^2 \zeta_{\mathbb{R}}(s - 2s_\infty) \zeta_{\mathbb{R}}(2s)^{-1} \\ &= L_\infty(s, \varrho_\infty \otimes \overline{\varrho}_\infty) \zeta_{\mathbb{R}}(2s)^{-1}. \end{aligned}$$

If $|k| \geq 2$,

$$\begin{aligned} Z_\infty(s, W_p^k \times \check{W}_p^k) &= (4\pi)^{-(s+|k|-1)} \Gamma(s + |k| - 1) \\ &= 2^{-|k|-1} \zeta_{\mathbb{C}}(s + |k| - 1) \zeta_{\mathbb{C}}(s) \zeta_{\mathbb{R}}(2s)^{-1} \\ &= 2^{-|k|-1} L_\infty(s, \varrho_\infty \otimes \overline{\varrho}_\infty) \zeta_{\mathbb{R}}(2s)^{-1}. \end{aligned}$$

If $p \nmid M$,

$$\begin{aligned} Z_p(s, W_p^0 \times \check{W}_p^0) &= \zeta_p(s + \dot{s}_p - \ddot{s}_p) \zeta_p(s)^2 \zeta_p(s + \ddot{s}_p - \dot{s}_p) \zeta_p(2s)^{-1} \\ &= L_p(s, \varrho_p \otimes \overline{\varrho}_p) \zeta_p(2s)^{-1}. \end{aligned}$$

If $p \mid M^\sharp$,

$$Z_p(s, W_p^\sharp \times \check{W}_p^\sharp) = \zeta_p(s + 1) = L_p(s, \varrho_p \otimes \overline{\varrho}_p) \zeta_p(s)^{-1}.$$

If $p \mid M^\chi$,

$$Z_p(s, W_p^\chi \times \check{W}_p^\chi) = \zeta_p(s) = L_p(s, \varrho_p \otimes \overline{\varrho}_p) \zeta_p(s)^{-1}.$$

By unfolding the integral of an incomplete Eisenstein series as above, one can show that $\text{res}_{s=1} E(s, g, \xi) = \frac{1}{2}$. Therefore the Petersson norm of F is given by

$$\begin{aligned} \langle F, F \rangle &= \frac{1}{2} \int_{PG_{\mathbb{Q}} \backslash PG_{\mathbb{A}}} |F(g)|^2 dg \\ &= \text{res}_{s=1} Z(s, F \times \overline{F}, \xi) \\ &= \lim_{s \rightarrow 1} Z(s, F \times \overline{F}, \xi) \zeta^*(s)^{-1} \\ &= \prod_v Z_v(1, \widehat{F}_v \times \widehat{\overline{F}}_v) \zeta_v(1)^{-1} \\ &= \int_{\mathbb{A}^\times} \left| \widehat{F} \left(\begin{bmatrix} a & \\ & 1 \end{bmatrix} \right) \right|^2 |a|_{\mathbb{A}}^{-1} da. \end{aligned}$$

It is easy to check the value of each c_v , defined by

$$Z_v(1, \widehat{F}_v \times \widehat{\overline{F}}_v) = \frac{c_v}{\zeta_v(2)} L_v(1, \varrho \otimes \overline{\varrho}).$$

The restriction $M^b = 1$ is removed using the calculation from §1.4,

$$\langle V_p^b, V_p^b \rangle = 2 \left(1 + \frac{\overline{\epsilon_p} \lambda_p}{p+1} \right) \langle \dot{V}_p^0, \dot{V}_p^0 \rangle.$$

□

Now we consider Eisenstein series which are Hecke eigenforms. For any choice of signs $\epsilon \in \prod_{p|M} \{\pm 1\}$, define $E(s, g, \xi^\epsilon)$ as before, with $R_p = R_p(1, (1, M))$ and

$$\xi^\epsilon(s, g) = \left(\prod_{p|M} (1 + \epsilon_p R_p) \right) \xi(s, g).$$

Then by Hejhal's calculation of the scattering matrix [10],

$$\begin{aligned} \eta(s, \xi^\epsilon) E(s, g, \xi^\epsilon) &= \eta(1-s, \xi^\epsilon) E(1-s, g, \xi^\epsilon), \\ \eta(s, \xi^\epsilon) &:= \left(\prod_{p|M} (1 + \epsilon_p p^s) \right) \zeta^*(2s). \end{aligned}$$

Finally, we make the spectral renormalization [10, 14],

$$\begin{aligned} E^1(s, g, \xi^\epsilon) &= \frac{2^{-\frac{1}{2}\#\{p|M\}+\frac{1}{2}}}{\text{vol}(\Gamma_0(M)\backslash\mathfrak{H})} E(s, g, \xi^\epsilon), \text{ so that} \\ \int_{\Gamma_0(M)\backslash\mathfrak{H}} \left| \int_0^\infty h(t) E^1\left(\frac{1}{2} + it, \sigma_z, \xi^\epsilon\right) \frac{dt}{\pi} \right|^2 \frac{dx dy}{y^2} &= \int_0^\infty |h(t)|^2 \frac{dt}{\pi}. \end{aligned}$$

Lemma 6. *Let $\psi \in S_k(1, (1, M), \chi, 1)$ be a classical Hecke normalized $\mathcal{H}^*$ eigenform, with Langlands parameters $\varrho = (\varrho_v)$. Then*

$$\frac{\int_{\Gamma_0(M)\backslash\mathfrak{H}} |\psi(z)|^2 E^1(s, \sigma_z, \xi^\epsilon) \frac{dx dy}{y^2}}{\int_{\Gamma_0(M)\backslash\mathfrak{H}} |\psi(z)|^2 \frac{dx dy}{y^2}} = \left(\prod_{p|M} \frac{1+\epsilon_p}{2} \right) \frac{2^{\frac{1}{2}\#\{p|M\}-\frac{3}{2}}}{M^{1-s}} \frac{L^*(s, \varrho \otimes \overline{\varrho})}{L^*(1, \text{Ad}\varrho) \eta(s, \xi^\epsilon)}.$$

Proof. Let F, f be as in Lemma 5, with $M^b = 1$. Since $R_p |\widehat{F}|^2 = |\widehat{F}|^2$ for all $p \mid M$,

$$\begin{aligned} Z(s, F \times \overline{F}, \xi^\epsilon) &= \left(\prod_{p|M} \frac{1+\epsilon_p}{2} \right) 2^{\#\{p|M\}} Z(s, F \times \overline{F}, \xi) \\ &= \left(\prod_{p|M} \frac{1+\epsilon_p}{2} \right) 2^{\#\{p|M\}} \frac{c_\infty L^*(s, \varrho \otimes \overline{\varrho})}{M^{-s} \eta(s, \xi^\epsilon)}, \text{ and} \\ Z(s, F \times \overline{F}, \xi^\epsilon) &= 2^{\frac{1}{2}\#\{p|M\}+\frac{1}{2}} \int_{\Gamma_0(M)\backslash\mathfrak{H}} |f(z)|^2 E^1(s, \sigma_z, \xi^\epsilon) \frac{dx dy}{y^2}. \end{aligned}$$

The result now follows by dividing out the similar formula of Lemma 5. Since the new formula is self-normalizing, we only require $\psi \in \mathbb{C}^\times f$ to be an eigenform. □

Note that the identity of Lemma 6 is consistent with the functional equations of $E^1(s, g, \xi^\epsilon)$ and $L^*(s, \varrho \otimes \overline{\varrho})$. For the purpose of comparison with Theorem 3, we observe that the Langlands parameters of the unitary Eisenstein series $E^1(\frac{1}{2} + it, g, \xi^\epsilon)$

are given by $\varrho_v^E = \|\cdot\|^{it} \oplus \|\cdot\|^{-it}$, and that

$$\frac{\left| \int_{\Gamma_0(M) \backslash \mathfrak{H}} |\psi(z)|^2 E^1\left(\frac{1}{2} + it, \sigma_z, \xi^\epsilon\right) \frac{dx dy}{y^2} \right|^2}{\left(\int_{\Gamma_0(M) \backslash \mathfrak{H}} |\psi(z)|^2 \frac{dx dy}{y^2} \right)^2} = (\prod_{p|M} Q_p^E) \frac{2^{\#\{p|M\}-3}}{M^2} \frac{L^*\left(\frac{1}{2}, \varrho \otimes \bar{\varrho} \otimes \varrho^E\right)}{L^*(1, \text{Ad}\varrho)^2 \text{res}_{s=1} L^*(s, \text{Ad}\varrho^E)},$$

$$Q_p^E = \left(\frac{1+\epsilon_p}{2}\right) (1 + p^{-\frac{1}{2}-it})^{-1} (1 + p^{-\frac{1}{2}+it})^{-1}.$$

3.2.2 Garrett/Piatetski-Shapiro–Rallis

In this section we explicitly calculate zeta integrals for the Rankin triple L-function, as defined in [33]. Let $F_j \in \pi_j \cap \tilde{S}_k(1, (1, d_B N^\sharp), 1, 1)$ be three Hecke normalized $\mathcal{H}_v^*$ eigenforms as in Theorem 1, with matching data φ_j as defined in §2.3. Consider $\mathbf{F} = F_1 \times F_2 \times F_3$ as an automorphic form on $PG_{\mathbb{A}}$ and $\boldsymbol{\varphi} = \varphi_1 \times \varphi_2 \times \varphi_3 \in \mathcal{S}(B_{\mathbb{A}}^3)$. We distinguish totally unramified data as $\boldsymbol{\varphi}^0$.

Following [33, 8], we consider $\Phi_v(s, h)$ satisfying

$$\Phi_v(s, th) = |\det AD^{-1}|_v^{s+1} \Phi_v(s, h), \quad \text{for } h \in H_v, \ t = \begin{bmatrix} A & B \\ 0 & D \end{bmatrix} \in T_v.$$

Note the Iwasawa decomposition $H_v = T_v K_v$, where

$$K_\infty = \text{U}(3), \quad K_p = \text{GSp}_6(\mathbb{Z}_p).$$

Now for $h \in H_v$, choose any $h' \in H'_v$ s.t. $\nu(h') = \nu(h)$ and define

$$\Phi_v(0, h) = \omega(h, h') \boldsymbol{\varphi}_v(0).$$

Since $\Phi_v^0(0, h)$ is constant on K_v ,

$$\Phi_v(s, h) = \left(\frac{\Phi_v^0(0, h)}{\Phi_v^0(0, 1)} \right)^s \Phi_v(0, h).$$

The local zeta integral is defined as in [33, 3, 7] by

$$Z_v(s, \widehat{\mathbf{F}}_v, \boldsymbol{\varphi}_v) = \int_{\mathbb{Q}_v^\times N_v^0 \backslash G_v} \Phi_v(s, \gamma_0 g) \widehat{\mathbf{F}}_v(g) dg,$$

where $N_v^0 = \left\{ \begin{bmatrix} & x_1 & & \\ & 1_3 & & \\ & & x_2 & \\ & & & x_3 \\ 0_3 & & & 1_3 \end{bmatrix} ; x_1 + x_2 + x_3 = 0 \right\},$

$$H_{\mathbb{Q}}^{(1)} \ni \gamma_0 = \begin{bmatrix} 1 & 1 & 1 & -1 & 0 & 0 \\ 0 & 1 & 0 & -1 & 1 & 0 \\ 0 & 0 & 1 & -1 & 0 & 1 \\ 1 & 1 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & -1 & 1 & 0 \\ 0 & 0 & 0 & -1 & 0 & 1 \end{bmatrix}.$$

Lemma 7. For any $\varphi_v \in \mathcal{S}(B_v^3)$,

$$\omega(\gamma_0)\varphi_v(0, 0, 0) = (-1)^{\mathbb{I}_v|d_B} \int_{B_v} \varphi_v(\beta, \beta, \beta) d\beta.$$

Proof. Write $\gamma_0 = J_1^{-1} E J_1 F G$,

$$E = \begin{bmatrix} & -1 & & \\ 1_3 & & 0 & \\ & & & 0 \\ 0_3 & & 1_3 & \end{bmatrix}, \quad J_1 = \begin{bmatrix} & 0 & & 1 \\ & 1_2 & & 0_2 \\ -1 & & 0 & \\ & 0_2 & & 1_2 \end{bmatrix},$$

$$F = \begin{bmatrix} & -1 & & \\ 1_3 & & 1 & \\ & & & 1 \\ 0_3 & & 1_3 & \end{bmatrix}, \quad G = \begin{bmatrix} 1 & 1 & 1 & & \\ & 1 & 0 & & \\ & & 1 & & \\ & & & 1 & \\ & & & -1 & 1 \\ & & & -1 & 0 & 1 \end{bmatrix}.$$

Then

$$\begin{aligned} \varphi'_v(\beta_1, \beta_2, \beta_3) &= \omega(FG)\varphi_v(\beta_1, \beta_2, \beta_3) \\ &= e_v(-\nu(\beta_1) + \nu(\beta_2) + \nu(\beta_3)) \\ &\quad \cdot \varphi_v(\beta_1, \beta_1 + \beta_2, \beta_1 + \beta_3), \end{aligned}$$

$$\begin{aligned} \omega(\gamma_0)\varphi_v(0, 0, 0) &= \omega(J_1^{-1} E J_1)\varphi'_v(0, 0, 0) \\ &= (\mathcal{F}_1^{-1} e_v(-\nu(\alpha_1)) \cdot \mathcal{F}_1 \varphi'_v)(0, 0, 0) \\ &= (-1)^{\mathbb{I}_v|d_B} \int_{B_v} e_v(\nu(\beta)) \varphi'_v(\beta, 0, 0) d\beta \\ &= (-1)^{\mathbb{I}_v|d_B} \int_{B_v} \varphi_v(\beta, \beta, \beta) d\beta. \end{aligned}$$

□

By the calculations of Piatetski-Shapiro–Rallis, Ikeda, Gross–Kudla, plus our own which follow, we have

Theorem 2. Suppose $k = |k_1| \geq |k_2| \geq |k_3| \geq 0$, $k_1 + k_2 + k_3 = 0$ ($\Rightarrow k_2 k_3 \geq 0$),

and for some fixed square-free N , all $N_j^\sharp = N$ and all $N_j^b = N_j^x = 1$. Then

$$Z_v(0, \check{\mathbf{W}}_v^\star, \boldsymbol{\varphi}_v) = \frac{C_v}{\zeta_v(2)^2} L_v(\tfrac{1}{2}, \pi_1 \times \pi_2 \times \pi_3),$$

$$C_\infty = \begin{cases} \frac{\varepsilon_\infty + 1}{2} & \text{if } k = 0, \\ 2^{-2k-2} & \text{if } k \geq 2. \end{cases} \quad C_p = \begin{cases} 1 & \text{if } p \nmid d_B N, \\ (\varepsilon_p - 1) \frac{p(p+\varepsilon_p)}{(p+1)^3} & \text{if } p \mid d_B, \\ (\varepsilon_p + 1) \frac{p(p+\varepsilon_p)}{(p+1)^3} & \text{if } p \mid N. \end{cases}$$

Proof. We go through each case separately. Note that although we only describe central values above, they are defined by analytic continuation of $Z_v(s)$.

Archimedean

Recall that

$$\kappa_\theta |F_j = e^{i|k_j|\theta} F_j, \quad \omega(\kappa_\theta) \varphi_j = e^{i|k_j|\theta} \varphi_j.$$

Defining

$$g_j = \begin{bmatrix} \epsilon a_j & \frac{x}{3} a_j^{-1} \\ & a_j^{-1} \end{bmatrix}, \quad h' = \rho \left(\begin{bmatrix} \epsilon & \\ & 1 \end{bmatrix}, 1 \right),$$

for $\epsilon \in \{\pm 1\}$, $a_j \in \mathbb{R}^\times$, $x \in \mathbb{R}$,

we have

$$\omega(g_j, h') \varphi_j(\beta) = \frac{1}{\pi^3} |a_j|^{k_j+2} X_\epsilon^{k_j}(\beta) e(\frac{x}{3}\nu(\beta) + i a_j^2 P(\beta)),$$

where $X_1 = X$, $X_{-1} = Y$. Then

$$\begin{aligned} \Phi(0, \delta g) &= \int_{M_2(\mathbb{R})} \omega(g) \boldsymbol{\varphi}(\beta, \beta, \beta) d\beta \\ &= \frac{1}{\pi^3} |a_1|^{k_1+2} |a_2|^{k_2+2} |a_3|^{k_3+2} \int_{M_2(\mathbb{R})} |X_\epsilon|^{2k} e(z|X|^2 - \bar{z}|Y|^2) d\beta, \\ &\quad \text{where } z = x + i(a_1^2 + a_2^2 + a_3^2) = x + iy, \\ &= \frac{1}{\pi^3} |a_1|^{k_1+2} |a_2|^{k_2+2} |a_3|^{k_3+2} \left[\mathcal{F} |X_\epsilon|^{2k} e(z|X|^2 - \bar{z}|Y|^2) \right] (0) \\ &= \frac{1}{\pi^3} (2\pi i)^{-2k} |a_1|^{k_1+2} |a_2|^{k_2+2} |a_3|^{k_3+2} \\ &\quad \cdot \left[\left(\frac{\partial}{\partial X_\epsilon} \frac{\partial}{\partial \bar{X}_\epsilon} \right)^k \mathcal{F} e(z|X|^2 - \bar{z}|Y|^2) \right] (0) \\ &= \frac{1}{\pi^3} (2\pi i)^{-2k} |a_1|^{k_1+2} |a_2|^{k_2+2} |a_3|^{k_3+2} |z|^{-2} \\ &\quad \cdot \left[\left(\frac{\partial}{\partial X_\epsilon} \frac{\partial}{\partial \bar{X}_\epsilon} \right)^k e\left(-\frac{1}{z}|X|^2 + \frac{1}{\bar{z}}|Y|^2\right) \right] (0) \\ &= \frac{1}{\pi^3} k! |a_1|^{k_1+2} |a_2|^{k_2+2} |a_3|^{k_3+2} (2\pi \epsilon z^\epsilon / i)^{-k} |z|^{-2}, \\ \Phi(s, \delta g) &= \frac{1}{\pi^3} k! |a_1|^{2s+k_1+2} |a_2|^{2s+k_2+2} |a_3|^{2s+k_3+2} (2\pi \epsilon z^\epsilon / i)^{-k} |z|^{-2s-2}. \end{aligned}$$

We used the formulas

$$\int_{\mathbb{R}^n} \exp(-\pi x^t Q x + 2\pi i y^t B x) dx = (\det Q)^{-\frac{1}{2}} \exp(-\pi y^t B Q^{-1} B^t y),$$

$$\left(\frac{\partial}{\partial z} \frac{\partial}{\partial \bar{z}}\right)^n \exp(a|z|^2) \Big|_{z=0} = n! a^n.$$

In the case of all F_j being even Maass forms, the difficult archimedean zeta integral was evaluated by Ikeda [12]. It is trivial to extend his result to arbitrary combinations of parity. Recall that

$$\{0, 1\} \ni \delta \equiv \delta_1 + \delta_2 + \delta_3 \pmod{2},$$

$$y = y_1 + y_2 + y_3.$$

Then

$$\begin{aligned} Z_\infty(s, \check{W}_\infty^0, \varphi_\infty) &= \sum_{\{\pm 1\}} \int_{(\mathbb{R}^\times)^3}^{\Pi > 0} \int_{\mathbb{R}} \int_{[0, \pi)^3} \frac{1}{\pi^3} |a_1 a_2 a_3|^{2s} |z|^{-2-2s} \\ &\quad \cdot \left(\prod_j \epsilon^{\delta_j} w_{0, s_j}(4\pi a_j^2) \right) e(x) 2 d^\times a_j dx d\theta_j \\ &= (1 - \delta) 2^4 \int_{(\mathbb{R}^+)^3} \int_{\mathbb{R}} |z|^{-2-2s} e(x) \prod_j a_j^{2s} w_{0, s_j}(4\pi a_j^2) d^\times a_j dx \\ &= (1 - \delta) 2^4 \int_{(\mathbb{R}^+)^3} \int_{\mathbb{R}} |z|^{-2-2s} e(x) \prod_j y_j^{s+\frac{1}{2}} K_{s_j}(2\pi y_j) d^\times y_j dx \\ &= \frac{(1 - \delta) 2^5 \pi^{s+1}}{\Gamma(s+1)} \int_{(\mathbb{R}^+)^3} y^{-s-\frac{1}{2}} K_{s+\frac{1}{2}}(2\pi y) \prod_j y_j^{s+\frac{1}{2}} K_{s_j}(2\pi y_j) d^\times y_j \\ &= \frac{(1 - \delta) \pi^{-s}}{\Gamma(s+1) \Gamma(2s+1)} \prod_{\{\pm\}^3} \Gamma\left(\frac{s}{2} + \frac{1}{4} \pm \frac{s_1}{2} \pm \frac{s_2}{2} \pm \frac{s_3}{2}\right) \\ &= \frac{(1 - \delta)}{\zeta_{\mathbb{R}}(2s+2) \zeta_{\mathbb{R}}(4s+2)} \prod_{\{\pm\}^3} \zeta_{\mathbb{R}}\left(s + \frac{1}{2} \pm s_1 \pm s_2 \pm s_3\right) \\ &= (1 - \delta) \frac{L_\infty\left(s + \frac{1}{2}, \pi_1 \times \pi_2 \times \pi_3\right)}{\zeta_{\mathbb{R}}(2s+2) \zeta_{\mathbb{R}}(4s+2)}. \end{aligned}$$

Our calculation in the case $|k_3| \geq 2$ is very similar to that of Gross–Kudla for all-equal positive weights [7].

$$\begin{aligned} Z_\infty(s, \check{W}_\infty^{|k_1|} \times \check{W}_\infty^{|k_2|} \times \check{W}_\infty^{|k_3|}, \varphi_\infty) &= \sum_{\{\pm 1\}} 4 \int_{(\mathbb{R}^+)^3} \mathbb{I}_{\mathbb{R}^-}(\epsilon) a_1^{2s+2|k_1|} a_2^{2s+2|k_2|} a_3^{2s+2|k_3|} \\ &\quad \cdot \int_{\mathbb{R}} k! (2\pi \epsilon z^\epsilon / i)^{-k} |z|^{-2s-2} e(z) 2 d^\times a_j dx \end{aligned}$$

$$\begin{aligned}
&= (2\pi i)^{-k} k! \int_{(\mathbb{R}^+)^3} y_1^{s+|k_1|} y_2^{s+|k_2|} y_3^{s+|k_3|} \\
&\quad \cdot \int_{\mathbb{R}} (\bar{z})^{-k} |z|^{-2s-2} e(z) d^\times y_j dx \\
&= (2\pi i)^{-k} k! \frac{(-2\pi i)^{s+1} (2\pi i)^{s+k+1}}{\Gamma(s+1) \Gamma(s+k+1)} \int_{(\mathbb{R}^+)^3} y_1^{s+|k_1|} y_2^{s+|k_2|} y_3^{s+|k_3|} \\
&\quad \cdot \int_0^\infty e^{-4\pi(y_1+y_2+y_3)(1+t)} t^s (1+t)^{s+k} d^\times y_j dt \\
&= \frac{(2\pi)^{2s+2} k!}{\Gamma(s+1) \Gamma(s+k+1)} \frac{\Gamma(s+|k_1|) \Gamma(s+|k_2|) \Gamma(s+|k_3|)}{(4\pi)^{3s+2k}} \\
&\quad \cdot \int_0^\infty t^s (1+t)^{-2s-k} dt \\
&= \frac{k! \Gamma(s+|k_1|) \Gamma(s+|k_2|) \Gamma(s+|k_3|)}{2^{4s+4k-2} \pi^{s+2k-2} \Gamma(s+1) \Gamma(s+k+1)} \frac{\Gamma(s+1) \Gamma(s+k-1)}{\Gamma(2s+k)} \\
&= \frac{2^{-2k-2} k!}{(s+k) (2s+1)_k} \frac{2^4 (2\pi)^{-4s-2k}}{\pi^{-3s-2}} \\
&\quad \cdot \frac{\Gamma(s+k-1) \Gamma(s+1) \Gamma(s+|k_2|) \Gamma(s+|k_3|)}{\Gamma(s+1) \Gamma(2s+1)} \\
&= \frac{2^{-2k-2} k!}{(s+k) (2s+1)_k} \frac{L_\infty(s+\frac{1}{2}, \pi_1 \times \pi_2 \times \pi_3)}{\zeta_{\mathbb{R}}(2s+2) \zeta_{\mathbb{R}}(4s+2)},
\end{aligned}$$

$$Z_\infty(0, \check{W}_\infty^{|k_1|} \times \check{W}_\infty^{|k_2|} \times \check{W}_\infty^{|k_3|}, \varphi_\infty) = \frac{2^{-2k-2}}{\zeta_{\mathbb{R}}(2)^2} L_\infty(\frac{1}{2}, \pi_1 \times \pi_2 \times \pi_3).$$

It is easy to check that our evaluation of the above integral is consistent with Ikeda's result for the overlapping parameter values $k_j = 0$, $s_j = -\frac{1}{2}$.

Finally, in the case $|k_2| \geq 2$, $|k_3| = 0$, we have

$$\begin{aligned}
&Z_\infty(s, \check{W}_\infty^k \times \check{W}_\infty^k \times \check{W}_\infty^0, \varphi_\infty) \\
&= (2\pi i)^{-k} k! \int_{(\mathbb{R}^+)^3} y_1^{s+k} y_2^{s+k} y_3^s w_{0,s_3}(4\pi y_3) e^{2\pi y_3} \\
&\quad \cdot \int_{\mathbb{R}} (\bar{z})^{-k} |z|^{-2s-2} e(z) d^\times y_j dx \\
&= (2\pi i)^{-k} k! \frac{(-2\pi i)^{s+1} (2\pi i)^{s+k+1}}{\Gamma(s+1) \Gamma(s+k+1)} \int_{(\mathbb{R}^+)^3} y_1^{s+k} y_2^{s+k} y_3^s w_{0,s_3}(4\pi y_3) \\
&\quad \cdot \int_0^\infty e^{-4\pi(y_1+y_2+y_3)(1+t)+2\pi y_3} t^s (1+t)^{s+k} d^\times y_j dt \\
&= \frac{(2\pi)^{2s+2} k!}{\Gamma(s+1) \Gamma(s+k+1)} \frac{\Gamma(s+k)^2}{(4\pi)^{2s+2k}} \int_{\mathbb{R}^+} y_3^s w_{0,s_3}(4\pi y_3) \\
&\quad \cdot \int_0^\infty e^{-2\pi y_3(1+2t)} t^s (1+t)^{-s-k} d^\times y_3 dt
\end{aligned}$$

$$\begin{aligned}
&= 2^{-2s-4k+2} \pi^{-2k+2} \frac{k! \Gamma(s+k)}{\Gamma(s+1)(s+k)} \frac{\Gamma(s+1)}{(4\pi)^s} \\
&\quad \cdot \int_{\mathbb{R}^+} (4\pi y_3)^{s+\frac{k}{2}-1} w_{-s-\frac{k}{2}, \frac{1-k}{2}}(4\pi y_3) w_{0,s_3}(4\pi y_3) d^\times y_3 \\
&= 2^{-4s-4k+2} \pi^{-s-2k+2} \frac{k! \Gamma(s+k)}{(s+k)} \\
&\quad \cdot \sum_{\{\pm\}^1} \frac{\Gamma(s+\frac{1}{2} \pm s_3) \Gamma(s+k-\frac{1}{2} \pm s_3) \Gamma(\mp 2s_3)}{\Gamma(\frac{1}{2} \mp s_3) \Gamma(2s+k+\frac{1}{2} \pm s_3)} \\
&\quad \cdot {}_3F_2 \left[\begin{matrix} s+\frac{1}{2} \pm s_3, & s+k-\frac{1}{2} \pm s_3, & \frac{1}{2} \pm s_3; \\ 1 \pm 2s_3, & 2s+k+\frac{1}{2} \pm s_3, & 1 \end{matrix} \right].
\end{aligned}$$

${}_3F_2$ denotes a generalized hypergeometric series. We can simplify this further at $s=0$ using the three-term relation,

$$\begin{aligned}
{}_3F_2 \left[\begin{matrix} a, & b, & c; \\ d, & e; & 1 \end{matrix} \right] &= \sum_{b \leftrightarrow c} \frac{\Gamma(1-a) \Gamma(d) \Gamma(e) \Gamma(c-b)}{\Gamma(d-b) \Gamma(e-b) \Gamma(1+b-a) \Gamma(c)} \\
&\quad \cdot {}_3F_2 \left[\begin{matrix} b, & 1+b-d, & 1+b-e; \\ 1+b-c, & 1+b-a; & 1 \end{matrix} \right].
\end{aligned}$$

$$\begin{aligned}
&Z_\infty(0, \check{W}_\infty^k \times \check{W}_\infty^k \times \check{W}_\infty^0, \varphi_\infty) \\
&= \frac{\Gamma(k)^2}{2^{4k-2} \pi^{2k-2}} \sum_{\{\pm\}^1} \frac{\Gamma(\frac{1}{2} \pm s_3) \Gamma(k-\frac{1}{2} \pm s_3) \Gamma(\mp 2s_3)}{\Gamma(\frac{1}{2} \mp s_3) \Gamma(k+\frac{1}{2} \pm s_3)} \\
&\quad \cdot {}_3F_2 \left[\begin{matrix} \frac{1}{2} \pm s_3, & k-\frac{1}{2} \pm s_3, & \frac{1}{2} \pm s_3; \\ 1 \pm 2s_3, & k+\frac{1}{2} \pm s_3; & 1 \end{matrix} \right] \\
&= \frac{\Gamma(k)}{2^{4k-2} \pi^{2k-2}} \sum_{\{\pm\}^1} \Gamma(\frac{1}{2} \mp s_3) \Gamma(-k+\frac{3}{2} \mp s_3) \Gamma(\frac{1}{2} \pm s_3) \Gamma(k-\frac{1}{2} \pm s_3) \\
&\quad \cdot \frac{\Gamma(k) \Gamma(\mp 2s_3)}{\Gamma(\frac{1}{2} \mp s_3) \Gamma(-k+\frac{3}{2} \mp s_3) \Gamma(\frac{1}{2} \mp s_3) \Gamma(k+\frac{1}{2} \pm s_3)} \\
&\quad \cdot {}_3F_2 \left[\begin{matrix} \frac{1}{2} \pm s_3, & k-\frac{1}{2} \pm s_3, & \frac{1}{2} \pm s_3; \\ 1 \pm 2s_3, & k+\frac{1}{2} \pm s_3; & 1 \end{matrix} \right] \\
&= \frac{\Gamma(k)}{2^{4k-2} \pi^{2k-2}} \frac{(-1)^{k-1} \pi^2}{\cos^2(\pi s_3)} \lim_{f \rightarrow 2-k} \frac{1}{\Gamma(f)} \\
&\quad \sum_{\{\pm\}^1} \frac{\Gamma(k) \Gamma(1) \Gamma(f) \Gamma(\mp 2s_3)}{\Gamma(\frac{1}{2} \mp s_3) \Gamma(f-\frac{1}{2} \mp s_3) \Gamma(k+\frac{1}{2} \pm s_3) \Gamma(\frac{1}{2} \mp s_3)} \\
&\quad \cdot {}_3F_2 \left[\begin{matrix} \frac{1}{2} \pm s_3, & \frac{1}{2} \pm s_3, & \frac{3}{2} \pm s_3 - f; \\ 1 \pm 2s_3, & k+\frac{1}{2} \pm s_3; & 1 \end{matrix} \right]
\end{aligned}$$

$$\begin{aligned}
&= \frac{\Gamma(k) (-1)^{k-1} \pi^2}{2^{4k-2} \pi^{2k-2} \cos^2(\pi s_3)} \lim_{f \rightarrow 2-k} \frac{1}{\Gamma(f)} {}_3F_2 \left[\begin{matrix} -k+1, & \frac{1}{2} + s_3, & \frac{1}{2} - s_3; \\ 1, & f; & 1 \end{matrix} \right] \\
&= \frac{\Gamma(k) (-1)^{k-1} \pi^2}{2^{4k-2} \pi^{2k-2} \cos^2(\pi s_3)} \sum_{n=k-1}^{\infty} \frac{(-k+1)_n (\frac{1}{2} + s_3)_n (\frac{1}{2} - s_3)_n}{n! (1)_n \Gamma(2-k+n)} \\
&= \frac{\Gamma(k) (-1)^{k-1} \pi^2}{2^{4k-2} \pi^{2k-2} \cos^2(\pi s_3)} \frac{(-1)^{k-1} \Gamma(k) \Gamma(k - \frac{1}{2} + s_3) \Gamma(k - \frac{1}{2} - s_3)}{\Gamma(k)^2 \Gamma(\frac{1}{2} + s_3) \Gamma(\frac{1}{2} - s_3)} \\
&= 2^{-4k+2} \pi^{-2k+2} \Gamma(k - \frac{1}{2} + s_3) \Gamma(k - \frac{1}{2} - s_3) \Gamma(\frac{1}{2} + s_3) \Gamma(\frac{1}{2} - s_3) \\
&= \frac{2^{-2k-2}}{\zeta_{\mathbb{R}}(2)^2} L_{\infty}(\frac{1}{2}, \pi_1 \times \pi_2 \times \pi_3).
\end{aligned}$$

This is consistent with the previous case if we set $s_3 = -\frac{1}{2}$. It would be interesting to directly factor the function $Z_{\infty}(s)$, for either $k = 0$ or $s_3 = -\frac{1}{2}$, by comparing with the previous cases.

Non-Archimedean

First consider $p \nmid d_B N$. A more illuminating treatment was given in [33], however our calculation is elementary and generalizes in an obvious way to the ramified cases. Note that the restriction $p \neq 3$ is unnecessary.

Writing $\alpha = \begin{bmatrix} a & \\ & 1 \end{bmatrix}$, $g_1 = \begin{bmatrix} a & x \\ & 1 \end{bmatrix}$, $g_j = \begin{bmatrix} aa_j & \\ & a_j^{-1} \end{bmatrix}$ for $j = 2, 3$, and choosing $h' = \rho(\alpha^t, 1)$, we compute

$$\begin{aligned}
\Phi_p(0, \gamma_0 g) &= \int_{M_2, \mathbb{Q}_p} \omega(g) \varphi_p(\beta, \beta, \beta) d\beta \\
&= |a^3 a_2^2 a_3^2|_p \int_{M_2, \mathbb{Q}_p} e_p(x\nu(\beta)) \varphi_{p1}(\alpha\beta) \varphi_{p2}(\alpha a_2 \beta) \varphi_{p3}(\alpha a_3 \beta) d\beta \\
&= \zeta_p(2)^3 |a^3 a_2^2 a_3^2|_p \int_{(\mathbb{Q}_p)^4} e_p(x(\beta_{11}\beta_{22} - \beta_{12}\beta_{21})) \\
&\quad \cdot \mathbb{I}_{\mathbb{Z}_p}(a\tilde{a}\beta_{11}) \mathbb{I}_{\dot{N}\mathbb{Z}_p}(a\tilde{a}\beta_{12}) \mathbb{I}_{\ddot{N}\mathbb{Z}_p}(\tilde{a}\beta_{21}) \mathbb{I}_{\mathbb{Z}_p}(\tilde{a}\beta_{22}) d\beta \\
&\quad (|\tilde{a}|_p := \max\{1, |a_2|_p, |a_3|_p\}) \\
&= \zeta_p(2)^3 |a^3 a_2^2 a_3^2|_p \int_{\mathbb{Q}_p} |a\tilde{a}|_p^{-1} \mathbb{I}_{a\tilde{a}\mathbb{Z}_p}(x\beta_{22}) \mathbb{I}_{\mathbb{Z}_p}(\tilde{a}\beta_{22}) d\beta_{22} \\
&\quad \cdot \int_{\mathbb{Q}_p} |\dot{N}|_p |a\tilde{a}|_p^{-1} \mathbb{I}_{a\tilde{a}\dot{N}\mathbb{Z}_p}(x\beta_{21}) \mathbb{I}_{\ddot{N}\mathbb{Z}_p}(\tilde{a}\beta_{21}) d\beta_{21} \\
&= \zeta_p(2)^3 |a^3 a_2^2 a_3^2|_p |z|_p^{-2} \\
&\quad (|z|_p := \max\{|a\tilde{a}^2|_p, |x|_p\}),
\end{aligned}$$

$$\Phi_p(s, \gamma_0 g) = \zeta_p(2)^3 |a^3 a_2^2 a_3^2|_p^{s+1} |z|_p^{-2s-2}.$$

It is straightforward to check that for $|a|_p \leq 1$,

$$\int_{\mathbb{Q}_p} |z|_p^{\lambda-1} e_p(x) dx = \left(\frac{1-p^{\lambda-1}}{1-p^{-\lambda}} \right) (1 - |pa\tilde{a}^2|_p^\lambda).$$

Then

$$\begin{aligned} Z_p(s, \check{\mathbf{W}}_p^0, \varphi_p) &= \int_{(\mathbb{Q}_p^\times)^3} \int_{\mathbb{Q}_p} |a^3 a_2^2 a_3^2|^{s+1} |z|^{-2s-2} e_p(x) \left(\frac{|ap|^{-\dot{s}_1} - |ap|^{-\dot{s}_1}}{|p|^{-\dot{s}_1} - |p|^{-\dot{s}_1}} \right) |a|^{\frac{1}{2}} \mathbb{I}_{\mathbb{Z}_p}(a) \\ &\quad \cdot \left(\frac{|aa_2 p|^{-\dot{s}_2} |a_2|^{\dot{s}_2} - |aa_2 p|^{-\dot{s}_2} |a_2|^{\dot{s}_2}}{|p|^{-\dot{s}_2} - |p|^{-\dot{s}_2}} \right) |aa_2^2|^{\frac{1}{2}} \mathbb{I}_{\mathbb{Z}_p}(aa_2^2) \\ &\quad \cdot \left(\frac{|aa_3 p|^{-\dot{s}_3} |a_3|^{\dot{s}_3} - |aa_3 p|^{-\dot{s}_3} |a_3|^{\dot{s}_3}}{|p|^{-\dot{s}_3} - |p|^{-\dot{s}_3}} \right) |aa_3^2|^{\frac{1}{2}} \mathbb{I}_{\mathbb{Z}_p}(aa_3^2) \\ &\quad \cdot |a^3 a_2^2 a_3^2|^{-1} dx d^\times a d^\times a_2 d^\times a_3 \\ &= \int_{|a| \leq 1} \int_{|a_2| \leq |a|^{-\frac{1}{2}}} \int_{|a_3| \leq |a|^{-\frac{1}{2}}} \left(\frac{1-p^{-2s-2}}{1-p^{2s+1}} \right) (1 - |pa\tilde{a}^2|^{-2s-1}) \\ &\quad \cdot \left(\frac{|ap|^{-\dot{s}_1} - |ap|^{-\dot{s}_1}}{|p|^{-\dot{s}_1} - |p|^{-\dot{s}_1}} \right) \left(\frac{|aa_2 p|^{-\dot{s}_2} |a_2|^{\dot{s}_2} - |aa_2 p|^{-\dot{s}_2} |a_2|^{\dot{s}_2}}{|p|^{-\dot{s}_2} - |p|^{-\dot{s}_2}} \right) \\ &\quad \cdot \left(\frac{|aa_3 p|^{-\dot{s}_3} |a_3|^{\dot{s}_3} - |aa_3 p|^{-\dot{s}_3} |a_3|^{\dot{s}_3}}{|p|^{-\dot{s}_3} - |p|^{-\dot{s}_3}} \right) |a^3 a_2^2 a_3^2|^{s+\frac{1}{2}} d^\times a d^\times a_2 d^\times a_3. \end{aligned}$$

Now write $|a|_p = p^{-k}$, $|a_j|_p = p^{-k_j}$, so

$$\begin{aligned} Z_p(s, \check{\mathbf{W}}_p^0, \varphi_p) &= \sum_{k \geq 0} \sum_{k_2 \geq -\frac{k}{2}} \sum_{k_3 \geq -\frac{k}{2}} S(k, k_2, k_3, \min\{k, k+2k_2, k+2k_3\}), \\ S(k, k_2, k_3, l) &:= \left(\frac{1-p^{-2s-2}}{1-p^{2s+1}} \right) p^{-(3k+2k_2+2k_3)(s+\frac{1}{2})} \\ &\quad \cdot \left(1 - p^{(l+1)(2s+1)} \right) \left(\frac{p^{(k+1)\dot{s}_1} - p^{(k+1)\dot{s}_1}}{p^{\dot{s}_1} - p^{\dot{s}_1}} \right) \\ &\quad \cdot \left(\frac{p^{(k+k_2+1)\dot{s}_2} - p^{(k+k_2+1)\dot{s}_2}}{p^{\dot{s}_2} - p^{\dot{s}_2}} \right) \\ &\quad \cdot \left(\frac{p^{(k+k_3+1)\dot{s}_3} - p^{(k+k_3+1)\dot{s}_3}}{p^{\dot{s}_3} - p^{\dot{s}_3}} \right). \end{aligned}$$

We can make the sums independent:

$$\begin{aligned} Z_p(s, \check{\mathbf{W}}_p^0, \varphi_p) &= \sum_{\mu=0,1} \sum_{k' \geq 0} \sum_{k'_2 \geq 0} \sum_{k'_3 \geq 0} \\ &\quad S(2k' + \mu, k'_2 - k', k'_3 - k', 2 \min\{k', k'_2, k'_3\} + \mu) \\ &= \sum_{\eta=0,1} \sum_{l \geq 0} \sum_{\mu=0,1} \sum_{k'' \geq 0} \sum_{k''_2 \geq 0} \sum_{k''_3 \geq 0} \\ &\quad (-1)^\eta S(2(k'' + l + \eta) + \mu, k''_2 - k'', k''_3 - k'', 2l + \mu). \end{aligned}$$

After multiplying out the summand as a polynomial and rearranging the exponents (as linear combinations of $\{1, l, k'', k''_2, k''_3\}$), we recognize Z_p to be a finite sum of products of geometric series. This evaluates to a complicated rational expression,

which we factor using Mathematica. It is necessary to impose the constraint on central characters, $\sum_j(\dot{s}_j + \ddot{s}_j) = 0$, e.g. by substitution for $\dot{s}_1$. Then

$$Z_p(s, \check{\mathbf{W}}_p^0, \boldsymbol{\varphi}_p) = \frac{L_p(s + \frac{1}{2}, \pi_1 \times \pi_2 \times \pi_3)}{\zeta_p(2s + 2) \zeta_p(4s + 2)}.$$

In the cases $p \mid d_B N^\sharp$, the difficult ramified zeta integrals were calculated by Gross–Kudla [7]. We have quoted their results, taking into account our differing normalizations of measures and local data. $\square$

Chapter 4

Conclusion

4.1 Triple Product Identities

Let $\psi_j \in S_{k_j}(d_B, \vec{N})$ be three Hecke-eigen newforms of the same square-free level N , with $k_1 + k_2 + k_3 = 0$, and let ϱ_j denote the corresponding Langlands parameters. Define $\varepsilon_v = \prod_j \varepsilon_{jv}$ for the eigenvalues ε_{jv} as in §1.4:

$$\begin{aligned} T_\infty^- \psi_j &= \varepsilon_{j\infty} \psi_j & \text{for } k_j = 0, \\ T_p^{[p]} \psi_j &= -\varepsilon_{jp} \psi_j & \text{for } p \mid d_B N. \end{aligned}$$

Writing $X = \mathcal{O}^{(1)}(d_B, \vec{N}) \backslash \mathfrak{H}$, we have

Theorem 3.

$$\frac{\left| \int_X \psi_1(z) \psi_2(z) \psi_3(z) \frac{dx dy}{y^2} \right|^2}{\prod_j \int_X |\psi_j(z)|^2 \frac{dx dy}{y^2}} = (\prod_v Q_v) \frac{2^{\#\{p \mid d_B N\} - 3} L^*(\frac{1}{2}, \varrho_1 \otimes \varrho_2 \otimes \varrho_3)}{(d_B N)^2} \frac{L^*(\frac{1}{2}, \varrho_1 \otimes \varrho_2 \otimes \varrho_3)}{\prod_j L^*(1, \text{Ad} \varrho_j)},$$

$$Q_\infty = \begin{cases} \frac{1+\varepsilon_\infty}{2} & \text{if } k_1 = k_2 = k_3 = 0, \\ 1 & \text{if } |k_1| = |k_2| > |k_3| = 0, \\ 2 & \text{if } |k_1| > |k_2| \geq |k_3| > 0, \end{cases} \quad Q_p = \begin{cases} \frac{1-\varepsilon_p}{2} & \text{if } p \mid d_B, \\ \frac{1+\varepsilon_p}{2} & \text{if } p \mid N, \\ 1 & \text{if } p \nmid d_B N. \end{cases}$$

Proof. Define the corresponding adelic forms Ψ_j on $B_\mathbb{A}^\times$, F_j on $G_\mathbb{A}$, F'_j on $H'_\mathbb{A}$, and the local data $\varphi_j \in \mathcal{S}(B_\mathbb{A})$, as in §1.2, §2.2. As Harris–Kudla showed, it follows from the see-saw identity,

$$\langle 1, \overline{\Theta_\varphi(\overline{\mathbf{F}})} \rangle_{PH'_\mathbb{A}} = \langle \Theta'_\varphi(1), \mathbf{F} \rangle_{PG_\mathbb{A}},$$

and the Siegel–Weil formula, $\Theta'_\varphi(1) = \frac{1}{2}E(0, h, \Phi)$, that

$$\int_{PH'_\mathbb{Q} \backslash PH'_\mathbb{A}} \Theta_\varphi(\overline{\mathbf{F}})(h') dh' = Z(0, \overline{\mathbf{F}}, \varphi).$$

Then by Theorem 1,

$$\frac{\left| \int_{PB_{\mathbb{Q}}^{\times} \setminus PB_{\mathbb{A}}^{\times}} \Psi_1(\beta) \Psi_2(\beta) \Psi_3(\beta) d^{\times} \beta \right|^2}{\prod_j \langle \Psi_j, \Psi_j \rangle_{PB_A^{\times}}} = \frac{Z(0, \overline{\mathbf{F}}, \boldsymbol{\varphi})}{\prod_j \langle F_j, F_j \rangle_{PG_{\mathbb{A}}}},$$

and by Theorem 2 and Lemma 5,

$$\frac{\left| \int_X \psi_1(z) \psi_2(z) \psi_3(z) \frac{dx dy}{y^2} \right|^2}{\prod_j \int_X |\psi_j(z)| \frac{dx dy}{y^2}} = \frac{\zeta^*(2)}{4 \text{vol}(X)} \frac{\prod_v C_v}{\prod_{j,v} c_{jv}} \frac{L^*(\frac{1}{2}, \varrho_1 \otimes \varrho_2 \otimes \varrho_3)}{\prod_j L^*(1, \text{Ad} \varrho_j)}.$$

Noting that

$$\text{vol}(X) = 2\zeta^*(2) \prod_{p|d_B} (p-1) \prod_{p|N} (p+1),$$

it is straightforward to compute the constants. $\square$

4.2 Applications to Quantum Chaos

Consider the geodesic flow on a finite volume hyperbolic surface $X = \Gamma \backslash \mathfrak{H}$. This is a classical Hamiltonian dynamical system with phase space T^*X , the cotangent bundle of X , and Hamiltonian $H(v_x^*) = |v_x^*|^2$, the Riemannian norm squared. We restrict our attention to S^*X , the constant energy submanifold of unit cotangent vectors, which is invariant under the flow. Under the homeomorphic identification

$$S^*X \xrightarrow{\sim} \Gamma \backslash \text{PGL}_2^+(\mathbb{R}),$$

$$(dy)_i \mapsto \Gamma \begin{bmatrix} 1 & \\ & 1 \end{bmatrix},$$

the flow map for time t is given explicitly by right multiplication,

$$T_t(\Gamma g) = \Gamma g \begin{bmatrix} e^t & \\ & 1 \end{bmatrix}.$$

The normalized Liouville measure on S^*X is ergodic for the geodesic flow and has entropy equal to 1. Furthermore, Ornstein–Weiss proved this system is measurably isomorphic to a Bernoulli flow. Topologically, the geodesic flow is less simple. There are no stable periodic orbits, but since the flow is Anosov, unstable periodic orbits form a dense subset. Each of these closed geodesics obviously carries an ergodic measure, but there are also ergodic measures with supports having any Hausdorff dimension between 1 and 3.

The standard quantization of the classical geodesic flow has state space $L^2(X)$ and Hamiltonian $\hat{H} = -\Delta$, a pseudo-differential operator with principal symbol H .

For a particle described by the normalized quantum state $\psi \in L^2(X)$, measuring its position is equivalent to sampling an X -valued random variable defined by the probability distribution

$$d\mu(z) = |\psi(z)|^2 \frac{dx dy}{y^2} \quad (\|\psi\|_{L^2} = 1).$$

Similarly, measuring the particle's energy is equivalent to sampling a $\text{Spec}(\tilde{H})$ -valued random variable, defined according to the spectral expansion of ψ . If X is compact, then $L^2(X)$ has an orthonormal basis of eigenforms,

$$\tilde{H}\psi_j = \lambda_j\psi_j, \quad 0 = \lambda_0 < \lambda_1 \leq \lambda_2 \leq \dots,$$

in which case the corresponding spectral measure is simply

$$\sum_j |\langle \psi, \psi_j \rangle|^2 \delta_{\lambda_j}.$$

For non-compact X , $\tilde{H}$ also has absolutely continuous spectrum equal to $[\frac{1}{4}, \infty)$. All of the corresponding (non-square-integrable) eigenspaces are spanned by unitary Eisenstein series and have dimension equal to the number of cusps of X .

The special arithmetic X we will consider are distinguished by their numerous symmetries, giving rise to Hecke operators. Since $\mathcal{H}^*$ is commutative, it represents a collection of simultaneously observable quantities. Furthermore, since $\tilde{H} \in \mathcal{H}_\infty^*$, we may assume that the ψ_j described above are in fact $\mathcal{H}^*$ -eigenforms. As we saw in Lemma 1 and the discussion before Lemma 6, $\mathcal{H}^*$ completely decomposes $L^2(X)$ with multiplicity 1. (Note that for these X , the residual spectrum consists of $\{0\}$, i.e. all ψ_j with $j > 0$ are cuspidal.)

Quantum chaos is concerned with the behavior of eigenstates in quantizations of classically chaotic systems, particularly under semi-classical/high energy limits [31]. An important general result due to Colin-de-Verdiere/Shnirelman/Zelditch is that

$$\frac{1}{N} \sum_{j=1}^N \left(\int_X f d\mu_j - \int_X f d\mu_0 \right)^2 \longrightarrow 0 \quad \text{as } N \rightarrow \infty, \text{ for all } f \in C_c^\infty(X).$$

This is called quantum ergodicity, and it is equivalent to the assertion that almost all high energy states are nearly equidistributed. The only assumption on X is ergodicity of the Liouville measure under geodesic flow, one of the mildest chaotic properties ensured by negative curvature. However, quantum ergodicity does not exclude the possibility that there may be other weak limits of the μ_j along subsequences, a phenomenon known as scarring. Scarring has been observed numerically in some related systems, such as the Bunimovich Stadium billiard. In the special cases of congruence hyperbolic surfaces (and 3-manifolds), Rudnick–Sarnak ruled out strong scarring along closed geodesics [29], leading them to conjecture that *all* high energy states become equidistributed. This is called quantum unique ergodicity (QUE). In [24], Luo–Sarnak made a quantitative formulation of the QUE conjecture, predicting

the rate of equidistribution: For all $f \in C_c^\infty(X)$,

$$\int_X f d\mu_j - \int_X f d\mu_0 \ll_{f,\epsilon} \lambda_j^{-\frac{1}{4}+\epsilon} \quad \text{as } \lambda_j \rightarrow \infty.$$

They proved this, in the case $X = \mathrm{SL}_2\mathbb{Z} \backslash \mathfrak{H}$, on average over μ_j and for the individual measures associated to unitary Eisenstein series. We now reduce their quantitative conjecture for individual μ_j to appropriate Lindelöf Hypotheses, using triple product identities. To quantify the smoothness of f , we use the Sobolev norms

$$\|f\|_{L^{2,r}} = \|(1 - \Delta)^{\frac{r}{2}} f\|_{L^2} \quad \text{for } r \in \mathbb{R}.$$

Let $L^{2,r}(X)$ denote the Hilbert space completion of $C_c^\infty(X)$ under $\|\cdot\|_{L^{2,r}}$.

Theorem 4. *Fix $X = \mathcal{O}^{(1)}(d_B) \backslash \mathfrak{H}$ and let the $\psi_j \in L^2(X)$ be as before, with Langlands parameters ϱ_j . Suppose that for some $0 \leq a \leq 1$ and all $j, j' \geq 1$,*

$$\begin{aligned} L(\tfrac{1}{2}, \varrho_j \otimes \overline{\varrho_j} \otimes \varrho_{j'}) &\ll_{X,\epsilon} C_\infty(0, \varrho_j \otimes \overline{\varrho_j} \otimes \varrho_{j'})^{\frac{a}{4}+\epsilon}, \quad \text{and also} \\ L(\tfrac{1}{2}, \varrho_j \otimes \overline{\varrho_j} \otimes \varrho_E) &\ll_{X,\epsilon} C_\infty(0, \varrho_j \otimes \overline{\varrho_j} \otimes \varrho_E)^{\frac{a}{4}+\epsilon} \quad \text{if } d_B = 1. \end{aligned}$$

Then for $0 \leq r \leq \frac{a+1}{2}$ and $\lambda_j \geq \frac{1}{4}$,

$$\left\| |\psi_j|^2 - \frac{1}{\mathrm{vol}(X)} \right\|_{L^{2,-r}} \ll_{X,\epsilon} \lambda_j^{\frac{a-r}{2}+\epsilon}.$$

Corollary 1. *Under the same hypotheses, set $r = \frac{a+1}{2}$. Then*

$$\int_X f d\mu_j - \int_X f d\mu_0 \ll_\epsilon \lambda_j^{\frac{a-1}{4}+\epsilon} \|f\|_{L^{2,r}}. \quad (4.1)$$

For $a = 1$, these hypotheses are Phragmen–Lindelöf convexity bounds, while an estimate quite similar to (4.1) follows trivially from the Sobolev inequality,

$$\|f\|_{L^\infty} \ll_{X,\epsilon} \|f\|_{L^{2,1+\epsilon}}.$$

Any value $a < 1$ (‘breaking convexity’) implies the QUE conjecture. Furthermore, the Grand Riemann Hypothesis implies $a = 0$, a generalization of Lindelöf’s Hypothesis, and this is best-possible. It follows from our proof that the corresponding exponent $-\frac{1}{4}$ of λ_j in (4.1) can’t be lowered. Our methods also apply to weight- k eigenforms ψ , giving the same result, although more smoothness is required from f . We will present these details later.

Corollary 2. *Under the same hypotheses, set $r = 0$. Then*

$$\|\psi_j\|_{L^4} \ll_\epsilon \lambda_j^{\frac{a}{4}+\epsilon}. \quad (4.2)$$

The exponent $\frac{1}{4}$ of λ_j in (4.2) corresponding to $a = 1$ is not the best which is presently known for general surfaces ($\frac{1}{16}$ is proved in [30]), and it is not the best

that can be done using Theorem 3. In joint work with Sarnak, we have obtained the exponent $\frac{1}{24}$ unconditionally, and we hope to improve this further.

Another related problem in quantum chaos concerns the amplitude distribution of high energy eigenstates. Viewing (X, μ_0) as a probability space, each ψ_j becomes an $\mathbb{R}$ -valued random variable with cumulative distribution function

$$f_j(t) = \mu_0(\psi_j^{-1}(-\infty, t]).$$

Berry/Hejhal's random wave conjecture asserts that these f_j converge in a suitable sense to the normal distribution $\mathcal{N}(0, \text{vol}(X)^{-\frac{1}{2}})$ as $\lambda_j \rightarrow \infty$. It follows from the proof of QE that the truth of the random wave conjecture for fourth moments alone would imply QUE. Regarding third moments, this conjecture says

$$\frac{1}{\text{vol}(X)} \int_X \psi_j^3(z) \frac{dx dy}{y^2} \longrightarrow 0 \quad \text{as } \lambda_j \rightarrow \infty.$$

We now prove a stronger form of this (unconditionally) for the full modular group.

Theorem 5. *Fix $X = \text{SL}_2\mathbb{Z} \backslash \mathfrak{H}$ and let the $\psi_j \in L^2(X)$ be defined as before. Then*

$$\int_X \psi_j^3(z) \frac{dx dy}{y^2} \ll_{\epsilon} \lambda_j^{-\frac{1}{12} + \epsilon}.$$

4.2.1 Proofs

Lemma 8. (Weyl's Law) *Writing $\lambda_j = \frac{1}{4} - s_j^2$, $s_j = \sigma_j + it_j$,*

$$\sum_{0 \leq t_j < T} 1 = \frac{\text{vol}(X)}{4\pi} T^2 + O_{X,\epsilon}((1+T)^{1+\epsilon}),$$

so for continuous f of bounded variation on $[a, b) \subset [0, \infty)$,

$$\begin{aligned} \sum_{a \leq t_j < b} f(t_j) - \frac{\text{vol}(X)}{2\pi} \int_a^b t f(t) dt \\ \ll_{X,\epsilon} (1+a)^{1+\epsilon} |f(a)| + (1+b)^{1+\epsilon} |f(b)| + \int_a^b (1+t)^{1+\epsilon} |df|(t). \end{aligned}$$

Proof of Theorem 4. Since $|\psi_j|^2 \in L^2(X)$, we can apply Parseval's formula:

$$\begin{aligned} \left\| |\psi_j|^2 - \frac{1}{\text{vol}(X)} \right\|_{L^2, -r}^2 &= \sum_{j' > 0} (1 + \lambda_{j'})^{-r} \left| \langle |\psi_j|^2, \psi_{j'} \rangle \right|^2 \\ &\quad + \int_{\mathbb{R}^+} \left(\frac{5}{4} + t^2 \right)^{-r} \left| \langle |\psi_j|^2, E^1(\cdot, \frac{1}{2} + it) \rangle \right|^2 \frac{dt}{\pi}, \end{aligned}$$

where the integral only appears if $d_B = 1$. First we analyze the individual terms, using Theorem 3 and Lemma 6. We may assume $\varepsilon_{j'\infty} = 1$, or else the corresponding

term vanishes by symmetry, and also $\lambda_j \geq \frac{1}{4}$. Then

$$\begin{aligned}
L_\infty(\tfrac{1}{2}, \varrho_j \times \overline{\varrho_j} \times \varrho_{j'}) &= \zeta_{\mathbb{R}}(\tfrac{1}{2} + 2s_j + s_{j'}) \zeta_{\mathbb{R}}(\tfrac{1}{2} + s_{j'})^2 \zeta_{\mathbb{R}}(\tfrac{1}{2} - 2s_j + s_{j'}) \\
&\quad \cdot \zeta_{\mathbb{R}}(\tfrac{1}{2} + 2s_j - s_{j'}) \zeta_{\mathbb{R}}(\tfrac{1}{2} - s_{j'})^2 \zeta_{\mathbb{R}}(\tfrac{1}{2} - 2s_j - s_{j'}), \\
C_\infty(0, \varrho_j \times \overline{\varrho_j} \times \varrho_{j'}) &= (1 + |2s_j + s_{j'}|)^2 (1 + |s_{j'}|)^4 (1 + |2s_j - s_{j'}|)^2, \\
L_\infty(\tfrac{1}{2}, \varrho_j \times \overline{\varrho_j} \times \varrho^E) &= \zeta_{\mathbb{R}}(\tfrac{1}{2} + 2s_j + it) \zeta_{\mathbb{R}}(\tfrac{1}{2} + it)^2 \zeta_{\mathbb{R}}(\tfrac{1}{2} - 2s_j + it) \\
&\quad \cdot \zeta_{\mathbb{R}}(\tfrac{1}{2} + 2s_j - it) \zeta_{\mathbb{R}}(\tfrac{1}{2} - it)^2 \zeta_{\mathbb{R}}(\tfrac{1}{2} - 2s_j - it), \\
C_\infty(0, \varrho_j \times \overline{\varrho_j} \times \varrho^E) &= (1 + |2s_j + it|)^2 (1 + |it|)^4 (1 + |2s_j - it|)^2.
\end{aligned}$$

Recall Stirling's approximation: For $s = \sigma + it$ and $|\arg(s)| \leq \pi - \epsilon < \pi$,

$$\begin{aligned}
\Gamma(s) &= (1 + O_\epsilon(|s|^{-1})) (2\pi)^{\frac{1}{2}} s^{s-\frac{1}{2}} e^{-s}, \\
|\Gamma(s)| &= (1 + O_\epsilon(|s|^{-1})) (2\pi)^{\frac{1}{2}} |s|^{\sigma-\frac{1}{2}} e^{-\sigma-t\arg(s)}.
\end{aligned}$$

We will write $X \asymp Y$ in place of $X \ll Y \ll X$. Then

$$\begin{aligned}
L_\infty(\tfrac{1}{2}, \varrho_j \times \overline{\varrho_j} \times \varrho_{j'}) &\asymp \frac{e^{-\frac{\pi}{2}(|2t_j+t_{j'}|+2|t_{j'}|+|2t_j-t_{j'}|)}}{C(0, \varrho_j \times \overline{\varrho_j} \times \varrho_{j'})^{\frac{1}{4}}}, \\
L_\infty(1, \text{Ad}\varrho_{j'}) &\asymp e^{-\pi|t_{j'}|}, \\
\frac{L_\infty(\tfrac{1}{2}, \varrho_j \times \overline{\varrho_j} \times \varrho_{j'})}{L_\infty(1, \text{Ad}\varrho_j)^2 L_\infty(1, \text{Ad}\varrho_{j'})} &\asymp \frac{e^{\frac{\pi}{2}(|2t_j+t_{j'}|-2|2t_j|+|2t_j-t_{j'}|)}}{C(0, \varrho_j \times \overline{\varrho_j} \times \varrho_{j'})^{\frac{1}{4}}}, \\
L_\infty(\tfrac{1}{2}, \varrho_j \times \overline{\varrho_j} \times \varrho^E) &\asymp \frac{e^{-\frac{\pi}{2}(|2t_j+t|+2|t|+|2t_j-t|)}}{C(0, \varrho_j \times \overline{\varrho_j} \times \varrho^E)^{\frac{1}{4}}}, \\
\text{res}_{z=1} L_\infty(z, \text{Ad}\varrho^E) &\asymp e^{-\pi|t|}, \\
\frac{L_\infty(\tfrac{1}{2}, \varrho_j \times \overline{\varrho_j} \times \varrho^E)}{L_\infty(1, \text{Ad}\varrho_j)^2 \text{res}_{z=1} L_\infty(z, \text{Ad}\varrho^E)} &\asymp \frac{e^{-\frac{\pi}{2}(|2t_j+t|-2|2t_j|+|2t_j-t|)}}{C(0, \varrho_j \times \overline{\varrho_j} \times \varrho^E)^{\frac{1}{4}}}.
\end{aligned}$$

Note that

$$|a+b| - 2|a| + |a-b| = 2 \max\{0, |b| - |a|\}.$$

Finally, recall the estimate of Hoffstein–Lockhart [11],

$$L(1, \text{Ad}\varrho_{j'})^{-1} \ll_\epsilon (1 + |t_{j'}|)^\epsilon \quad \text{for all } \epsilon > 0.$$

We divide the discrete spectrum contributions into

$$\text{I} = \{0 \leq t_{j'} < t_j\}, \quad \text{II} = \{t_j \leq t_{j'} < 3t_j\}, \quad \text{III} = \{3t_j \leq t_{j'}\}.$$

Then

$$\begin{aligned}
\sum_I &\ll_\epsilon (1+t_j)^{a-1+\epsilon} \sum_I (1+t_{j'})^{a-1-2r+\epsilon} \\
&\ll_\epsilon (1+t_j)^{\max\{a-1, 2a-2r\}+\epsilon} \ll_\epsilon (1+t_j)^{2a-2r+\epsilon}, \\
\sum_I I &\ll_\epsilon (1+t_j)^{\frac{3}{2}(a-1)-2r+\epsilon} \sum_I I (1+|2t_j-t_{j'}|)^{\frac{1}{2}(a-1)+\epsilon} \\
&\ll_\epsilon (1+t_j)^{2a-2r+\epsilon}, \\
\sum_I II &\ll_\epsilon (1+t_j)^\epsilon \sum_I II (1+t_{j'})^{2a-2-2r+\epsilon} e^{-\pi(t_{j'}-2t_j)} \\
&\ll_\epsilon e^{-(\pi-\epsilon)t_j} \ll (1+t_j)^{2a-2r}.
\end{aligned}$$

Similarly, we divide up the continuous spectrum:

$$\begin{aligned}
\int_0^{t_j} &\ll_\epsilon (1+t_j)^{a-1+\epsilon} \int_0^{t_j} (1+t)^{a-1-2r+\epsilon} dt \\
&\ll_\epsilon (1+t_j)^{\max\{a-1, 2a-2r-1\}+\epsilon} \ll_\epsilon (1+t_j)^{2a-2r+\epsilon}, \\
\int_{t_j}^{3t_j} &\ll_\epsilon (1+t_j)^{\frac{3}{2}(a-1)-2r+\epsilon} \int_{t_j}^{3t_j} (1+|2t_j-t|)^{\frac{1}{2}(a-1)+\epsilon} dt \\
&\ll_\epsilon (1+t_j)^{2a-2r-1+\epsilon}, \\
\int_{3t_j}^\infty &\ll_\epsilon (1+t_j)^\epsilon \int_{3t_j}^\infty (1+t)^{2a-2-2r+\epsilon} e^{-\pi(t-2t_j)} dt \\
&\ll_\epsilon e^{-(\pi-\epsilon)t_j} \ll (1+t_j)^{2a-2r}.
\end{aligned}$$

□

Proof of Theorem 5. First note that

$$L_v(s, \otimes^3 \varrho_j) = L_v(s, \text{Sym}^3 \varrho_j) L_v(s, \varrho_j)^2.$$

Then for $\lambda_j \geq \frac{1}{4}$,

$$\frac{L_\infty(\frac{1}{2}, \otimes^3 \varrho_j)}{L_\infty(\frac{1}{2}, \text{Ad} \varrho_j)^3} \asymp (1+|t_j|)^{-2}, \quad C_\infty(0, \text{Sym}^3 \varrho_j) \asymp (1+|t_j|)^4.$$

To prove the convexity bound for $L(s, \text{Sym}^3 \varrho_j)$, we first note that it is entire by the work of Kim–Shahidi [19], and has finite order by the work of Gelbart–Shahidi [5]. Furthermore, using the converse theorem of Cogdell–Piatetski-Shapiro [2], Kim–Shahidi proved that $\text{Sym}^3 \varrho_j$ is cuspidal on GL_4 [20]. Recent results of Molteni [25]

then imply

$$\begin{aligned} L(1+it, \text{Sym}^3 \varrho_j) &\ll_\epsilon C_\infty(t, \text{Sym}^3 \varrho_j)^\epsilon, \\ L(0+it, \text{Sym}^3 \varrho_j) &\ll_\epsilon C_\infty(t, \text{Sym}^3 \varrho_j)^{\frac{1}{2}+\epsilon}, \end{aligned}$$

and hence by the Phragmen–Lindelöf/Hadamard three-circles method,

$$L(\tfrac{1}{2}, \text{Sym}^3 \varrho_j) \ll_\epsilon C(0, \text{Sym}^3 \varrho_j)^{\frac{1}{4}+\epsilon} \ll (1+|t_j|)^{1+\epsilon}.$$

Now to prove that the third moment tends to zero, we only need subconvexity for $L(\frac{1}{2}, \varrho_j)$. The first such estimate was proved conditionally by Iwaniec in [14]; his result is made unconditional at a small cost to the exponent by the same technique as in [15]. The current record bound is due to Ivić and subsequently Jutila [13, 18]:

$$L(\tfrac{1}{2}, \varrho_j) \ll_\epsilon (1+|t_j|)^{\frac{1}{3}+\epsilon}.$$

Therefore

$$\left| \int_X \psi_j^3(z) \frac{dx dy}{y^2} \right|^2 \ll_\epsilon (1+|t_j|)^{-\frac{1}{3}+\epsilon}.$$

□

Bibliography

- [1] S. Böcherer, R. Schulze-Pillot, On the central critical value of the triple product L -function, *Number theory* (Paris, 1993–1994), London Math. Soc. Lecture Note Ser. 235, Cambridge Univ. Press, Cambridge (1996), 1–46.
- [2] J. Cogdell, I. Piatetski-Shapiro, A converse theorem for GL_4 , *Math. Res. Lett.* 3 (1996), 67–76.
- [3] P. Garrett, M. Harris, Special values of triple product L -functions, *Amer. J. Math.* 115 (1993), 161–240.
- [4] S. Gelbart, *Automorphic Forms on Adele Groups*, Annals of Mathematics Studies 83, Princeton Univ. Press (1975)
- [5] S. Gelbart, F. Shahidi, Boundedness of automorphic L -functions in vertical strips, *J. Amer. Math. Soc.* 14 (2001), 79–107.
- [6] R. Godement, *Notes on Jacquet–Langlands Theory*, IAS mimeograph (1970)
- [7] B. Gross and S. Kudla, Heights and the central critical values of triple product L -functions, *Compositio Math.* 81 (1992), 143–209.
- [8] M. Harris and S. Kudla, The central critical value of a triple product L -function, *Annals of Math.* 133 (1991), 605–702
- [9] M. Harris and S. Kudla, Arithmetic automorphic forms for the non-holomorphic discrete series of $GS(2)$, *Duke Math. J.* 66 (1992), 59–121.
- [10] D.A. Hejhal, *The Selberg Trace Formula for $PSL(2, R)$* , Lecture Notes in Math. 1001 (1983)
- [11] J. Hoffstein, P. Lockhart, Coefficients of Maass forms and the Siegel zero, *Ann. of Math.* (2) 140 (1994), 161–181.
- [12] T. Ikeda, On the gamma factor of the triple L -function, I, *Duke Math. J.* 97 (1999), 301–318.
- [13] A. Ivić, On sums of Hecke series in short intervals, *to appear*
- [14] H. Iwaniec, The spectral growth of automorphic L -functions, *J. Reine Angew. Math.* 428 (1992), 139–159.

- [15] H. Iwaniec, P. Sarnak, L^∞ norms of eigenfunctions of arithmetic surfaces, *Ann. of Math.* (2) 141 (1995), 301–320.
- [16] H. Iwaniec, P. Sarnak, Perspectives on the analytic theory of L -functions, GAFA 2000 (Tel Aviv, 1999), *Geom. Funct. Anal.* (2000) Special Volume, Part II, 705–741.
- [17] H. Jacquet, R. Langlands, *Automorphic forms on $GL(2)$* , Lecture Notes in Math. 114, Springer (1970)
- [18] M. Jutila, The fourth moment of central values of Hecke series, *Number theory* (Turku, 1999), de Gruyter, Berlin (2001), 167–177.
- [19] H. Kim, F. Shahidi, Symmetric cube L -functions for GL_2 are entire, *Ann. of Math.* 150 (1999), 645–662.
- [20] H. Kim, F. Shahidi, Functorial products for $GL_2 \times GL_3$ and functorial symmetric cube for GL_2 , *C. R. Acad. Sci. Paris Sr. I Math.* 331 (2000), 599–604.
- [21] A. Knapp, Local Langlands correspondence: the Archimedean case, *Motives* (Seattle, 1991), Proc. Sympos. Pure Math. 55, Part 2, Amer. Math. Soc., Providence, RI (1994), 393–410.
- [22] S. Kudla, The local Langlands correspondence: the non-Archimedean case, *Motives* (Seattle, 1991), Proc. Sympos. Pure Math. 55, Part 2, Amer. Math. Soc., Providence, RI (1994), 365–391.
- [23] G. Lione and M. Vergne, *The Weil representation, Maslov index, and theta series*, Birkhauser, Boston (1980)
- [24] W. Luo, P. Sarnak, Quantum ergodicity of eigenfunctions on $PSL_2(\mathbf{Z}) \backslash \mathbf{H}^2$, *Inst. Hautes Études Sci. Publ. Math.* 81 (1995), 207–237.
- [25] G. Molteni, Upper and lower bounds for certain L -functions at $s = 1$ for certain Dirichlet series with Euler product, *to appear in Duke Math. J.*
- [26] T. Miyake, On automorphic forms on GL_2 and Hecke operators, *Ann. of Math.* 94 (1971), 174–189.
- [27] T. Miyake, *Modular Forms*, Springer–Verlag, Berlin (1989)
- [28] A. Pizer, Hecke operators for $\Gamma_0(N)$, *J. Algebra* 83 (1983), 39–64.
- [29] Z. Rudnick, P. Sarnak, The behaviour of eigenstates of arithmetic hyperbolic manifolds, *Comm. Math. Phys.* 161 (1994), 195–213.
- [30] A. Seeger, C. Sogge, Bounds for eigenfunctions of differential operators, *Indiana Univ. Math. J.* 38 (1989), 669–682.
- [31] P. Sarnak, Arithmetic quantum chaos, The Schur lectures (Tel Aviv, 1992), *Israel Math. Conf. Proc.* 8 (1995), 183–236.

- [32] F. Shahidi, Local coefficients as Artin factors for real groups, *Duke Math. J.* 52 (1985), 973–1007.
- [33] I. Piatetski-Shapiro and S. Rallis, Rankin triple L -functions, *Comp. Math.* 64 (1987), 31–115.
- [34] H. Shimizu, Theta series and automorphic forms on GL_2 , *J. Math. Soc. Japan* 24 (1972), 638–683.
- [35] H. Shimizu, On zeta functions of quaternion algebras, *Ann. of Math.* 81 (1965), 166–193.
- [36] G. Shimura, *Introduction to the Arithmetic Theory of Automorphic Functions*, Princeton University Press, Princeton (1971)
- [37] G. Shimura, On certain zeta functions attached to two Hilbert modular forms. II. The case of automorphic forms on a quaternion algebra, *Ann. of Math.* 114 (1981), 569–607.
- [38] D. Ornstein, B. Weiss, Geodesic flows are Bernoullian, *Israel J. Math.* 14 (1973), 184–198.
- [39] J. Tate, Number theoretic background, *Automorphic forms, representations and L -functions* (Corvallis, 1977), Proc. Sympos. Pure Math. 33, Part 2, Amer. Math. Soc., Providence, R.I. (1979), 3–26.
- [40] M-F. Vignéras, *Arithmétique des algèbres de quaternions*, Lecture Notes in Math. 800, Springer, Berlin (1980)
- [41] A. Weil, *Basic Number Theory*, Die Grundlehren der mathematischen Wissenschaften, Band 144 Springer-Verlag, New York (1967)
- [42] A. Weil, *Adeles and Algebraic Groups*, Progress in Mathematics 23, Birkhuser, Boston (1982)